

# Max-flow and min-cut problems

Max Flow and  
Min Cut

Properties of  
flows and cuts

Residual  
graph

Augmenting  
path

MaxFlow  
MinCut Thm

Ford  
Fulkerson alg

Maximum  
matching in  
Bip graphs

Disjoint paths  
problem



# 1 Max Flow and Min Cut

2 Properties of flows and cuts

3 Residual graph

4 Augmenting path

5 MaxFlow MinCut Thm

6 Ford Fulkerson alg

7 Maximum matching in Bip graphs

Max Flow and  
Min Cut

Properties of  
flows and cuts

Residual  
graph

Augmenting  
path

MaxFlow  
MinCut Thm

Ford  
Fulkerson alg

Maximum  
matching in  
Bip graphs

Disjoint paths  
problem

# Flow Network

## Max Flow and Min Cut

Properties of flows and cuts

Residual graph

Augmenting path

MaxFlow MinCut Thm


Ford Fulkerson alg

Maximum matching in Bip graphs

Disjoint paths problem

A network  $\mathcal{N} = (V, E, c, s, t)$  is formed by

- a digraph  $G = (V, E)$ ,
- a source vertex  $s \in V$
- a sink vertex  $t \in V$ ,
- and edge capacities  $c : E \rightarrow \mathbb{R}^+$



# A flow in a network


Given a network  $\mathcal{N} = (V, E, c, s, t)$

A **Flow** is an assignment  $f : E \rightarrow \mathbb{R}^+ \cup \{0\}$  that follows the **Kirchoff's laws**:

- $\forall (u, v) \in E, 0 \leq f(u, v) \leq c(u, v),$
- (Flow conservation)  $\forall v \in V - \{s, t\},$   
 $\sum_{u \in V} f(u, v) = \sum_{z \in V} f(v, z)$

The **value of a flow**  $f$  is

$$|f| = \sum_{v \in V} f(s, v) = f(s, V) = f(V, t).$$



$$f(e)/c(e)$$

Max Flow and  
Min Cut

Properties of  
flows and cuts

Residual  
graph

Augmenting  
path

MaxFlow  
MinCut Thm

Ford  
Fulkerson alg

Maximum  
matching in  
Bip graphs

Disjoint paths  
problem

# A flow in a network


Given a network  $\mathcal{N} = (V, E, c, s, t)$

A **Flow** is an assignment  $f : E \rightarrow \mathbb{R}^+ \cup \{0\}$  that follows the **Kirchoff's laws**:

- $\forall (u, v) \in E, 0 \leq f(u, v) \leq c(u, v),$
- (Flow conservation)  $\forall v \in V - \{s, t\},$   
$$\sum_{u \in V} f(u, v) = \sum_{z \in V} f(v, z)$$

The **value of a flow**  $f$  is

$$|f| = \sum_{v \in V} f(s, v) = f(s, V) = f(V, t).$$



$$f(e)/c(e)$$

with value 3.

Max Flow and  
Min Cut

Properties of  
flows and cuts

Residual  
graph

Augmenting  
path

MaxFlow  
MinCut Thm

Ford  
Fulkerson alg

Maximum  
matching in  
Bip graphs

Disjoint paths  
problem

# A flow in a network


Given a network  $\mathcal{N} = (V, E, c, s, t)$

A **Flow** is an assignment  $f : E \rightarrow \mathbb{R}^+ \cup \{0\}$  that follows the **Kirchoff's laws**:

- $\forall (u, v) \in E, 0 \leq f(u, v) \leq c(u, v),$
- (Flow conservation)  $\forall v \in V - \{s, t\},$   
$$\sum_{u \in V} f(u, v) = \sum_{z \in V} f(v, z)$$

The **value of a flow**  $f$  is

$$|f| = \sum_{v \in V} f(s, v) = f(s, V) = f(V, t).$$



saturated

Max Flow and  
Min Cut

Properties of  
flows and cuts

Residual  
graph

Augmenting  
path

MaxFlow  
MinCut Thm

Ford  
Fulkerson alg

Maximum  
matching in  
Bip graphs

Disjoint paths  
problem

# The Maximum flow problem

INPUT: A network  $\mathcal{N} = (V, E, c, s, t,)$

QUESTION: Find a flow of maximum value on  $\mathcal{N}$ .

Max Flow and  
Min Cut

Properties of  
flows and cuts

Residual  
graph

Augmenting  
path

MaxFlow  
MinCut Thm

Ford  
Fulkerson alg


Maximum  
matching in  
Bip graphs

Disjoint paths  
problem

# The Maximum flow problem

INPUT: A network  $\mathcal{N} = (V, E, c, s, t,)$

QUESTION: Find a flow of maximum value on  $\mathcal{N}$ .



Max Flow and  
Min Cut

Properties of  
flows and cuts

Residual  
graph

Augmenting  
path

MaxFlow  
MinCut Thm

Ford  
Fulkerson alg

Maximum  
matching in  
Bip graphs


Disjoint paths  
problem



# The Maximum flow problem

INPUT: A network  $\mathcal{N} = (V, E, c, s, t,)$

QUESTION: Find a flow of maximum value on  $\mathcal{N}$ .



The value of the flow is  $7 = 4 + 1 + 2 = 5 + 2$ .

As  $t$  cannot receive more flow, this flow is a **maximum flow**.

Max Flow and  
Min Cut

Properties of  
flows and cuts

Residual  
graph

Augmenting  
path

MaxFlow  
MinCut Thm

Ford  
Fulkerson alg

Maximum  
matching in  
Bip graphs


Disjoint paths  
problem

# The $(s, t)$ -cuts

Given  $\mathcal{N} = (V, E, c, s, t)$  a  $(s, t)$ -cut is a partition of  $V = S \cup T$  ( $S \cap T = \emptyset$ ), with  $s \in S$  and  $t \in T$ .

The capacity of a cut  $(S, T)$  is the sum of weights leaving  $S$ , i.e.,

$$c(S, T) = \sum_{u \in S} \sum_{v \in T} c(u, v)$$



$$S = \{s, c, d\}$$

$$T = \{a, b, e, t\}$$

$$c(S, T) = 19$$

$$(4 + 5) + 5 + (3 + 2)$$

Max Flow and Min Cut

Properties of flows and cuts

Residual graph

Augmenting path

MaxFlow MinCut Thm

Ford Fulkerson alg

Maximum matching in Bip graphs


Disjoint paths problem

# The $(s, t)$ -cuts

Given  $\mathcal{N} = (V, E, c, s, t)$  a  $(s, t)$ -cut is a partition of  $V = S \cup T$  ( $S \cap T = \emptyset$ ), with  $s \in S$  and  $t \in T$ .

The flow across the cut:

$$f(S, T) = \sum_{u \in S} \sum_{v \in T} f(u, v) - \sum_{v \in T} \sum_{u \in S} f(v, u).$$



$$\begin{aligned} S &= \{s, c, d\} \\ T &= \{a, b, e, t\} \\ c(S, T) &= 19 \\ f(S, T) &= 10 - 3 = 7 \end{aligned}$$

Max Flow and Min Cut

Properties of flows and cuts

Residual graph

Augmenting path

MaxFlow MinCut Thm

Ford Fulkerson alg

Maximum matching in Bip graphs


Disjoint paths problem

# The $(s, t)$ -cuts

Given  $\mathcal{N} = (V, E, c, s, t)$  a  $(s, t)$ -cut is a partition of  $V = S \cup T$  ( $S \cap T = \emptyset$ ), with  $s \in S$  and  $t \in T$ .

The flow across the cut:

$$f(S, T) = \sum_{u \in S} \sum_{v \in T} f(u, v) - \sum_{v \in T} \sum_{u \in S} f(v, u).$$



$$\begin{aligned} S &= \{s, c, d\} \\ T &= \{a, b, e, t\} \\ c(S, T) &= 19 \\ f(S, T) &= 10 - 3 = 7 \end{aligned}$$

Due to the capacity constrain:  $f(S, T) \leq c(S, T)$

Max Flow and Min Cut

Properties of flows and cuts

Residual graph

Augmenting path

MaxFlow MinCut Thm

Ford Fulkerson alg

Maximum matching in Bip graphs


Disjoint paths problem

## Another $(s, t)$ -cut

Given  $\mathcal{N} = (V, E, c, s, t)$  a  $(s, t)$ -cut is a partition of  $V = S \cup T$  ( $S \cap T = \emptyset$ ), with  $s \in S$  and  $t \in T$ .

The flow across the cut:

$$f(S, T) = \sum_{u \in S} \sum_{v \in T} f(u, v) - \sum_{v \in T} \sum_{u \in S} f(v, u).$$



$$S = \{s, a, c\}$$

$$T = \{b, d, e, t\}$$

$$c(S, T) = 12$$

$$f(S, T) = 8 - 1 = 7$$

Max Flow and  
Min Cut

Properties of  
flows and cuts

Residual  
graph

Augmenting  
path

MaxFlow  
MinCut Thm

Ford  
Fulkerson alg

Maximum  
matching in  
Bip graphs

Disjoint paths  
problem

# The Minimum Cut problem

Max Flow and  
Min Cut

Properties of  
flows and cuts

Residual  
graph

Augmenting  
path

MaxFlow  
MinCut Thm


Ford  
Fulkerson alg

Maximum  
matching in  
Bip graphs

Disjoint paths  
problem

INPUT: A network  $\mathcal{N} = (V, E, c, s, t,)$

QUESTION: Find a  $(s, t)$ -cut of minimum capacity in  $\mathcal{N}$ .



MinCut

$S = \{s, a, b, c, d, e\}$

$T = \{t\}$

$c(S, T) = 7$

# Changing weights effect on min cuts

Given a network  $\mathcal{N} = (V, E, s, t, c)$  assume that  $(S, T)$  is a min  $(s, t)$ -cut.

Max Flow and  
Min Cut

Properties of  
flows and cuts

Residual  
graph

Augmenting  
path

MaxFlow  
MinCut Thm

Ford  
Fulkerson alg



Maximum  
matching in  
Bip graphs

Disjoint paths  
problem

# Changing weights effect on min cuts

Given a network  $\mathcal{N} = (V, E, s, t, c)$  assume that  $(S, T)$  is a min  $(s, t)$ -cut.

If we change the input by adding  $c > 0$  to the capacity of every edge, then it may happen that  $(S, T)$  is not longer a min  $(s, t)$ -cut.



Max Flow and Min Cut

Properties of flows and cuts

Residual graph

Augmenting path

MaxFlow MinCut Thm

Ford Fulkerson alg

Maximum matching in Bip graphs

Disjoint paths problem



# Changing weights effect on Min-Cut and Max-Flow

Given a network  $\mathcal{N} = (V, E, s, t, c)$ .

Max Flow and  
Min Cut

Properties of  
flows and cuts

Residual  
graph

Augmenting  
path

MaxFlow  
MinCut Thm

Ford  
Fulkerson alg

Maximum  
matching in  
Bip graphs

Disjoint paths  
problem

# Changing weights effect on Min-Cut and Max-Flow

Max Flow and  
Min Cut

Properties of  
flows and cuts

Residual  
graph

Augmenting  
path

MaxFlow  
MinCut Thm



Ford  
Fulkerson alg

Maximum  
matching in  
Bip graphs

Disjoint paths  
problem

Given a network  $\mathcal{N} = (V, E, s, t, c)$ .

If we change the network by multiplying by  $c >$  the capacity of every edge, the capacity of any  $(s, t)$ -cut in the new network is  $c$  times its capacity in the original network.



1 Max Flow and Min Cut

2 Properties of flows and cuts

3 Residual graph

4 Augmenting path

5 MaxFlow MinCut Thm

6 Ford Fulkerson alg

7 Maximum matching in Bip graphs

Max Flow and  
Min Cut

Properties of  
flows and cuts

Residual  
graph

Augmenting  
path

MaxFlow  
MinCut Thm

Ford  
Fulkerson alg

Maximum  
matching in  
Bip graphs

Disjoint paths  
problem

# Notation

Let  $\mathcal{N} = (V, E, s, t, c)$  and  $f$  a flow in  $\mathcal{N}$

For  $v \in V$ ,  $U \subseteq V$  and  $v \notin U$ .

- $f(v, U)$  flow  $v \rightarrow U$  i.e.  $f(v, U) = \sum_{u \in U} f(v, u)$ ,
- $f(U, v)$  flow  $U \rightarrow v$  i.e.  $f(U, v) = \sum_{u \in U} f(u, v)$ ,

For a  $(s, t)$ -cut  $(S, T)$  and  $v \in S$

- $S' = S \setminus \{v\}$  and  $T' = T \cup \{v\}$
- $f_{-v}(S, T) = \sum_{u \in S'} \sum_{w \in T} f(u, w) - \sum_{w \in T} \sum_{u \in S'} f(w, u)$   
i.e, the contribution to  $f(S, T)$  from edges not incident with  $v$ .

Max Flow and  
Min Cut

Properties of  
flows and cuts

Residual  
graph

Augmenting  
path

MaxFlow  
MinCut Thm

Ford  
Fulkerson alg

Maximum  
matching in  
Bip graphs

Disjoint paths  
problem

# Flow conservation on $(s, t)$ -cuts

Max Flow and  
Min Cut

Properties of  
flows and cuts

Residual  
graph

Augmenting  
path

MaxFlow  
MinCut Thm

Ford  
Fulkerson alg

Maximum  
matching in  
Bip graphs

Disjoint paths  
problem

## Theorem

*Let  $\mathcal{N} = (V, E, s, t, c)$  and  $f$  a flow in  $\mathcal{N}$ . For any  $(s, t)$ -cut  $(S, T)$ ,  $f(S, T) = |f|$ .*

**Proof** (Induction on  $|S|$ )

- If  $S = \{s\}$  then, by definition,  $f(S, T) = |f|$ .

# Flow conservation on $(s, t)$ -cuts

Max Flow and  
Min Cut

Properties of  
flows and cuts

Residual  
graph

Augmenting  
path

MaxFlow  
MinCut Thm

Ford  
Fulkerson alg

Maximum  
matching in  
Bip graphs

Disjoint paths  
problem

## Theorem


Let  $\mathcal{N} = (V, E, s, t, c)$  and  $f$  a flow in  $\mathcal{N}$ . For any  $(s, t)$ -cut  $(S, T)$ ,  $f(S, T) = |f|$ .

## Proof (Induction on $|S|$ )

- If  $S = \{s\}$  then, by definition,  $f(S, T) = |f|$ .
- Assume it is true for  $S' = S - \{v\}$  and  $T' = T \cup \{v\}$ , i.e.  $f(S', T') = |f|$ .

# Flow conservation on $(s, t)$ -cuts

## Proof (cont.) (Induction on $|S|$ )



- IH:  $f(S', T') = |f|$ .
- Then,  $f(S, T) = f_{-v}(S, T) + f(v, T) - f(T, v)$ .
- But,  $f(S', T') = f_{-v}(S, T) + f(S', v) - f(v, S')$  as  $v \in T'$
- By flow conservation,  
 $f(S', v) + f(T, v) = f(v, S') + f(v, T)$
- So,  $f(S', v) - f(v, S') = f(v, T) - f(T, v)$
- Therefore,  $f(S', T') = f(S, T) = |f|$

- 1 Max Flow and Min Cut
- 2 Properties of flows and cuts
- 3 Residual graph**
- 4 Augmenting path
- 5 MaxFlow MinCut Thm
- 6 Ford Fulkerson alg
- 7 Maximum matching in Bip graphs

Max Flow and  
Min Cut

Properties of  
flows and cuts

**Residual  
graph**

Augmenting  
path

MaxFlow  
MinCut Thm

Ford  
Fulkerson alg

Maximum  
matching in  
Bip graphs



Disjoint paths  
problem



# Residual graph

Given a network  $\mathcal{N} = (V, E, s, t, c)$  together with a **flow**  $f$ .  
The **residual graph**,  $(G_f = (V, E_f, c_f))$  is a weighted digraph on the same vertex set and with edge set:

- if  $c(u, v) - f(u, v) > 0$ , then  $(u, v) \in E_f$  and  $c_f(u, v) = c(u, v) - f(u, v) > 0$  (**forward edges**)



Max Flow and  
Min Cut

Properties of  
flows and cuts

**Residual  
graph**

Augmenting  
path

MaxFlow  
MinCut Thm

Ford  
Fulkerson alg



Maximum  
matching in  
Bip graphs

Disjoint paths  
problem

# Residual graph

Given a network  $\mathcal{N} = (V, E, s, t, c)$  together with a flow  $f$  on it, the residual graph,  $(G_f = (V, E_f, c_f))$  is a weighted digraph on the same vertex set and with edge set:

- if  $f(u, v) > 0$ , then  $(v, u) \in E_f$  and  $c_f(v, u) = f(u, v)$  (**backward edges**).



Max Flow and Min Cut

Properties of flows and cuts

Residual graph

Augmenting path

MaxFlow MinCut Thm

Ford Fulkerson alg



Maximum matching in Bip graphs

Disjoint paths problem

# Residual graph

Given a network  $\mathcal{N} = (V, E, s, t, c)$  together with a flow  $f$  on it, the residual graph,  $(G_f = (V, E_f, c_f))$  is a weighted digraph on the same vertex set and with edge set:

- if  $c(u, v) - f(u, v) > 0$ , then  $(u, v) \in E_f$  and  $c_f(u, v) = c(u, v) - f(u, v) > 0$  (forward edges)
- if  $f(u, v) > 0$ , then  $(v, u) \in E_f$  and  $c_f(v, u) = f(u, v)$  (backward edges).



Max Flow and Min Cut

Properties of flows and cuts

Residual graph

Augmenting path



MaxFlow MinCut Thm

Ford Fulkerson alg

Maximum matching in Bip graphs

Disjoint paths problem

# Residual graph



- Notice that, if  $c(u, v) = f(u, v)$ , then there is only a backward edge.
- $c_f$  are called the residual capacity.

Max Flow and  
Min Cut

Properties of  
flows and cuts

Residual  
graph

Augmenting  
path



MaxFlow  
MinCut Thm

Ford  
Fulkerson alg

Maximum  
matching in  
Bip graphs

Disjoint paths  
problem

# Residual graph



- **forward edges:** There remains capacity to push more flow through this edge.
- **backward edges:** there are units of flow that can be redirected through other links.

Max Flow and  
Min Cut

Properties of  
flows and cuts

**Residual  
graph**

Augmenting  
path

MaxFlow  
MinCut Thm

Ford  
Fulkerson alg

Maximum  
matching in  
Bip graphs

Disjoint paths  
problem

1 Max Flow and Min Cut

2 Properties of flows and cuts

3 Residual graph

**4 Augmenting path**

5 MaxFlow MinCut Thm

6 Ford Fulkerson alg

7 Maximum matching in Bip graphs

Max Flow and  
Min Cut

Properties of  
flows and cuts

Residual  
graph

**Augmenting  
path**

MaxFlow  
MinCut Thm



Ford  
Fulkerson alg

Maximum  
matching in  
Bip graphs

Disjoint paths  
problem

# Augmenting paths

Let  $\mathcal{N} = (V, E, c, s, t)$  and let  $f$  be a flow in  $\mathcal{N}$ ,



- An **augmenting path**  $P$  is any **simple** path  $P$  in  $G_f$  from  $s$  to  $t$

Max Flow and  
Min Cut

Properties of  
flows and cuts

Residual  
graph

**Augmenting  
path**

MaxFlow  
MinCut Thm



Ford  
Fulkerson alg

Maximum  
matching in  
Bip graphs

Disjoint paths  
problem

# Augmenting paths

Let  $\mathcal{N} = (V, E, c, s, t)$  and let  $f$  be a flow in  $\mathcal{N}$ ,



- An **augmenting path**  $P$  is any **simple path**  $P$  in  $G_f$  from  $s$  to  $t$ .  $P$  might have forward and backward edges.

Max Flow and  
Min Cut

Properties of  
flows and cuts

Residual  
graph

**Augmenting  
path**

MaxFlow  
MinCut Thm

Ford  
Fulkerson alg



Maximum  
matching in  
Bip graphs

Disjoint paths  
problem



# Augmenting paths

Let  $\mathcal{N} = (V, E, c, s, t)$  and let  $f$  be a flow in  $\mathcal{N}$ ,



- An **augmenting path**  $P$  is any **simple** path  $P$  in  $G_f$  from  $s$  to  $t$ .  $P$  might have forward and backward edges.
- For an augmenting path  $P$  in  $G_f$ , the **bottleneck**,  $b(P)$ , is the minimum (residual) capacity of the edges in  $P$ . In the example, for  $P = (s, b, a, t)$ ,  $b(P) = 1$ .

Max Flow and  
Min Cut

Properties of  
flows and cuts

Residual  
graph

**Augmenting  
path**

MaxFlow  
MinCut Thm


Ford  
Fulkerson alg

Maximum  
matching in  
Bip graphs

Disjoint paths  
problem

# Augmenting paths: increasing the flow

```
Augment( $P, f$ )  
   $b$  = bottleneck ( $P$ )  
  for each  $(u, v) \in P$  do  
    if  $(u, v)$  is a forward edge then  
      Increase  $f(u, v)$  by  $b$   
    else  
      Decrease  $f(v, u)$  by  $b$   
  return  $f$ 
```



Max Flow and  
Min Cut

Properties of  
flows and cuts

Residual  
graph

**Augmenting  
path**

MaxFlow  
MinCut Thm



Ford  
Fulkerson alg

Maximum  
matching in  
Bip graphs

Disjoint paths  
problem

# Augmenting paths: increasing the flow

**Augment**( $P, f$ )  
 $b = \text{bottleneck}(P)$   
**for each**  $(u, v) \in P$  **do**  
    **if**  $(u, v)$  is a forward edge **then**  
        Increase  $f(u, v)$  by  $b$   
    **else**  
        Decrease  $f(v, u)$  by  $b$   
**return**  $f$



$G_f, P = (s, a, t), b(P) = 1$

Max Flow and  
Min Cut

Properties of  
flows and cuts

Residual  
graph

Augmenting  
path

MaxFlow  
MinCut Thm




Ford  
Fulkerson alg

Maximum  
matching in  
Bip graphs

Disjoint paths  
problem

# Augmenting paths: increasing the flow

**Augment**( $P, f$ )  
 $b$ =bottleneck ( $P$ )  
**for each**  $(u, v) \in P$  **do**  
    **if**  $(u, v)$  is a forward edge **then**  
        Increase  $f(u, v)$  by  $b$   
    **else**  
        Decrease  $f(v, u)$  by  $b$   
**return**  $f$



Max Flow and  
Min Cut

Properties of  
flows and cuts

Residual  
graph

Augmenting  
path

MaxFlow  
MinCut Thm

Ford  
Fulkerson alg

Maximum  
matching in  
Bip graphs

Disjoint paths  
problem

# Augmenting paths: increasing the flow

Max Flow and  
Min Cut

Properties of  
flows and cuts

Residual  
graph

**Augmenting  
path**

MaxFlow  
MinCut Thm

Ford  
Fulkerson alg

Maximum  
matching in  
Bip graphs

Disjoint paths  
problem

## Lemma

*Let  $f' = \text{Augment}(P, f)$ , then  $f'$  is a flow in  $G$ .*

**Proof:** We have to prove the two flow properties.

### ■ Capacity law

- Forward edges  $(u, v) \in P$ , we increase  $f(u, v)$  by  $b$ , as  $b \leq c(u, v) - f(u, v)$  then  $f'(u, v) = f(u, v) + b \leq c(u, v)$ .
- Backward edges  $(u, v) \in P$  we decrease  $f(v, u)$  by  $b$ , as  $b \leq f(v, u)$ ,  $f'(v, u) = f(v, u) - b \geq 0$ .

# Augmenting paths: increasing the flow

## Lemma

Let  $f' = \text{Augment}(P, f)$ , then  $f'$  is a flow in  $G$ .

**Proof:** We have to prove the two flow properties.

- **Conservation law**,  $\forall v \in P \setminus \{s, t\}$  let  $u$  be the predecessor of  $v$  in  $P$  and let  $w$  be its successor.
- As the path is simple only the alterations due to  $(u, v)$  and  $(v, w)$  can change the flow that goes through  $v$ . We have four cases:
  - $(u, v)$  and  $(v, w)$  are backward edges, the flow in  $(v, u)$  and  $(w, v)$  is decremented by  $b$ . As one is incoming and the other outgoing the total balance is 0.
  - $(u, v)$  and  $(v, w)$  are forward edges, the flow in  $(u, v)$  and  $(v, w)$  is incremented by  $b$ . As one is incoming and the other outgoing the total balance is 0.

# Augmenting paths: increasing the flow

## Lemma

Let  $f' = \text{Augment}(P, f)$ , then  $f'$  is a flow in  $G$ .

**Proof:** We have to prove the two flow properties.

- **Conservation law**,  $\forall v \in P \setminus \{s, t\}$  let  $u$  be the predecessor of  $v$  in  $P$  and let  $w$  be its successor.
- As the path is simple only the alterations due to  $(u, v)$  and  $(v, w)$  can change the flow that goes through  $v$ . We have three cases:
  - $(u, v)$  is forward and  $(v, w)$  is backward, the flow in  $(u, v)$  is incremented by  $b$  and the flow in  $(w, v)$  is decremented by  $b$ . As both are incoming, the total balance is 0.
  - $(u, v)$  is backward and  $(v, w)$  is forward, the flow in  $(v, w)$  is incremented by  $b$  and the flow in  $(v, u)$  is decremented by  $b$ . As both are outgoing, the total balance is 0.

# Augmenting paths: incrementing the flow

Max Flow and  
Min Cut

Properties of  
flows and cuts

Residual  
graph

**Augmenting  
path**

MaxFlow  
MinCut Thm

Ford  
Fulkerson alg

Maximum  
matching in  
Bip graphs

Disjoint paths  
problem

## Lemma

*Consider  $f' = \text{Augment}(P, f)$ , then  $|f'| > |f|$ .*

**Proof:** Let  $P$  be the augmenting path in  $G_f$ . The first edge  $e \in P$  leaves  $s$ , and as  $G$  has no incoming edges to  $s$ ,  $e$  is a forward edge. Moreover  $P$  is simple  $\Rightarrow$  never returns to  $s$ . Therefore, the value of the flow increases in edge  $e$  by  $b$  units.

□



- 1 Max Flow and Min Cut
- 2 Properties of flows and cuts
- 3 Residual graph
- 4 Augmenting path
- 5 MaxFlow MinCut Thm**
- 6 Ford Fulkerson alg
- 7 Maximum matching in Bip graphs

Max Flow and  
Min Cut

Properties of  
flows and cuts

Residual  
graph

Augmenting  
path

**MaxFlow  
MinCut Thm**

Ford  
Fulkerson alg

Maximum  
matching in  
Bip graphs

Disjoint paths  
problem

# Max-Flow Min-Cut theorem

Ford and Fulkerson (1954); Peter Elias, Amiel Feinstein and Claude Shannon (1956) (in framework of information-theory).

## Theorem

*For any  $\mathcal{N}(G, s, t, c)$ , the maximum of the flow value is equal to the minimum of the  $(S, T)$ -cut capacities.*

$$\max_f \{ |f| \} = \min_{(S, T)} \{ c(S, T) \}.$$

Max Flow and  
Min Cut

Properties of  
flows and cuts

Residual  
graph

Augmenting  
path

**MaxFlow  
MinCut Thm**

Ford  
Fulkerson alg

Maximum  
matching in  
Bip graphs

Disjoint paths  
problem

# Max-Flow Min-Cut theorem: Proof

## Proof:

- Let  $f^*$  be a flow with maximum value,  $|f^*| = \max_f \{|f|\}$

Max Flow and  
Min Cut

Properties of  
flows and cuts

Residual  
graph

Augmenting  
path

**MaxFlow  
MinCut Thm**

Ford  
Fulkerson alg

Maximum  
matching in  
Bip graphs

Disjoint paths  
problem

# Max-Flow Min-Cut theorem: Proof

## Proof:

- Let  $f^*$  be a flow with maximum value,  $|f^*| = \max_f \{|f|\}$
- For any  $(s, t)$ -cut  $(S, T)$ ,  $f^*(S, T) \leq c(S, T)$ .

Max Flow and  
Min Cut

Properties of  
flows and cuts

Residual  
graph

Augmenting  
path

**MaxFlow  
MinCut Thm**

Ford  
Fulkerson alg

Maximum  
matching in  
Bip graphs

Disjoint paths  
problem

# Max-Flow Min-Cut theorem: Proof

## Proof:

- Let  $f^*$  be a flow with maximum value,  $|f^*| = \max_f \{|f|\}$
- For any  $(s, t)$ -cut  $(S, T)$ ,  $f^*(S, T) \leq c(S, T)$ .
- $G_{f^*}$  has no augmenting path. So, if  $S_s = \{v \in V \mid \exists s \rightsquigarrow v \text{ in } G_{f^*}\}$ , then  $(S_s, V - \{S_s\})$  is a  $(s, t)$ -cut.

Max Flow and  
Min Cut

Properties of  
flows and cuts

Residual  
graph

Augmenting  
path

MaxFlow  
MinCut Thm

Ford  
Fulkerson alg

Maximum  
matching in  
Bip graphs

Disjoint paths  
problem

# Max-Flow Min-Cut theorem: Proof

## Proof:

- Let  $f^*$  be a flow with maximum value,  $|f^*| = \max_f \{|f|\}$
- For any  $(s, t)$ -cut  $(S, T)$ ,  $f^*(S, T) \leq c(S, T)$ .
- $G_{f^*}$  has no augmenting path. So, if  $S_s = \{v \in V \mid \exists s \rightsquigarrow v \text{ in } G_{f^*}\}$ , then  $(S_s, V - \{S_s\})$  is a  $(s, t)$ -cut.
- For  $e = (u, v) \in E$  with  $u \in S_s$  and  $v \notin S_s$ ,  $(u, v) \notin E(G_{f^*})$ , therefore  $f^*(u, v) = c(u, v)$ ,

Max Flow and  
Min Cut

Properties of  
flows and cuts

Residual  
graph

Augmenting  
path

MaxFlow  
MinCut Thm

Ford  
Fulkerson alg

Maximum  
matching in  
Bip graphs

Disjoint paths  
problem

# Max-Flow Min-Cut theorem: Proof

## Proof:

- Let  $f^*$  be a flow with maximum value,  $|f^*| = \max_f \{|f|\}$
- For any  $(s, t)$ -cut  $(S, T)$ ,  $f^*(S, T) \leq c(S, T)$ .
- $G_{f^*}$  has no augmenting path. So, if  $S_s = \{v \in V \mid \exists s \rightsquigarrow v \text{ in } G_{f^*}\}$ , then  $(S_s, V - \{S_s\})$  is a  $(s, t)$ -cut.
- For  $e = (u, v) \in E$  with  $u \in S_s$  and  $v \notin S_s$ ,  $(u, v) \notin E(G_{f^*})$ , therefore  $f^*(u, v) = c(u, v)$ ,
- Then,  $c(S_s, V - \{S_s\}) = f^*(S_s, V - \{S_s\}) = |f^*|$

Max Flow and  
Min Cut

Properties of  
flows and cuts

Residual  
graph

Augmenting  
path

MaxFlow  
MinCut Thm

Ford  
Fulkerson alg

Maximum  
matching in  
Bip graphs

Disjoint paths  
problem

# Max-Flow Min-Cut theorem: Proof

## Proof:

- Let  $f^*$  be a flow with maximum value,  $|f^*| = \max_f \{|f|\}$
- For any  $(s, t)$ -cut  $(S, T)$ ,  $f^*(S, T) \leq c(S, T)$ .
- $G_{f^*}$  has no augmenting path. So, if  $S_s = \{v \in V \mid \exists s \rightsquigarrow v \text{ in } G_{f^*}\}$ , then  $(S_s, V - \{S_s\})$  is a  $(s, t)$ -cut.
- For  $e = (u, v) \in E$  with  $u \in S_s$  and  $v \notin S_s$ ,  $(u, v) \notin E(G_{f^*})$ , therefore  $f^*(u, v) = c(u, v)$ ,
- Then,  $c(S_s, V - \{S_s\}) = f^*(S_s, V - \{S_s\}) = |f^*|$
- $(S_s, V - \{S_s\})$  is a minimum capacity  $(s, t)$ -cut in  $G$ .



Max Flow and  
Min Cut

Properties of  
flows and cuts

Residual  
graph

Augmenting  
path

MaxFlow  
MinCut Thm

Ford  
Fulkerson alg

Maximum  
matching in  
Bip graphs

Disjoint paths  
problem



1 Max Flow and Min Cut

2 Properties of flows and cuts

3 Residual graph

4 Augmenting path

5 MaxFlow MinCut Thm

**6 Ford Fulkerson alg**

7 Maximum matching in Bip graphs

Max Flow and  
Min Cut

Properties of  
flows and cuts

Residual  
graph

Augmenting  
path

MaxFlow  
MinCut Thm


**Ford  
Fulkerson alg**

Maximum  
matching in  
Bip graphs

Disjoint paths  
problem

# Ford-Fulkerson algorithm

L.R. Ford, D.R. Fulkerson:  
*Maximal flow through a network*. Canadian J. of Math. 1956.



```
Ford-Fulkerson( $G, s, t, c$ )  
for all  $(u, v) \in E$  set  $f(u, v) = 0$   
 $G_f = G$   
while there is an  $(s, t)$  path  $P$  in  $G_f$  do  
     $f = \text{Augment}(P, G_f)$   
    Compute  $G_f$   
return  $f$ 
```

Max Flow and  
Min Cut

Properties of  
flows and cuts

Residual  
graph

Augmenting  
path


MaxFlow  
MinCut Thm

Ford  
Fulkerson alg

Maximum  
matching in  
Bip graphs

Disjoint paths  
problem

# FF algorithm example



Max Flow and  
Min Cut

Properties of  
flows and cuts

Residual  
graph

Augmenting  
path



MaxFlow  
MinCut Thm

**Ford  
Fulkerson alg**

Maximum  
matching in  
Bip graphs

Disjoint paths  
problem

# FF algorithm example



Max Flow and  
Min Cut

Properties of  
flows and cuts

Residual  
graph

Augmenting  
path


MaxFlow  
MinCut Thm

**Ford  
Fulkerson alg**


Maximum  
matching in  
Bip graphs

Disjoint paths  
problem


# FF algorithm example



Initial  $f_0$



$G_{f_0}$



$f_1$

Max Flow and  
Min Cut

Properties of  
flows and cuts

Residual  
graph

Augmenting  
path


MaxFlow  
MinCut Thm

Ford  
Fulkerson alg


Maximum  
matching in  
Bip graphs

Disjoint paths  
problem


# FF algorithm example




Initial  $f_0$



$G_{f_0}$



$f_1$



$G_{f_1}$

Max Flow and Min Cut

Properties of flows and cuts

Residual graph

Augmenting path



MaxFlow MinCut Thm

Ford Fulkerson alg

Maximum matching in Bip graphs

Disjoint paths problem

# FF example



Max Flow and Min Cut

Properties of flows and cuts

Residual graph

Augmenting path




MaxFlow  
MinCut Thm

**Ford  
Fulkerson alg**

Maximum  
matching in  
Bip graphs

Disjoint paths  
problem

# FF example



Max Flow and  
Min Cut

Properties of  
flows and cuts

Residual  
graph

Augmenting  
path

MaxFlow  
MinCut Thm





Ford  
Fulkerson alg

Maximum  
matching in  
Bip graphs

Disjoint paths  
problem



# FF example



Max Flow and Min Cut

Properties of flows and cuts

Residual graph

Augmenting path



MaxFlow MinCut Thm

Ford Fulkerson alg

Maximum matching in Bip graphs

Disjoint paths problem

# FF example



Max Flow and  
Min Cut

Properties of  
flows and cuts

Residual  
graph

Augmenting  
path




MaxFlow  
MinCut Thm

**Ford  
Fulkerson alg**

Maximum  
matching in  
Bip graphs

Disjoint paths  
problem

# FF example



Max Flow and Min Cut

Properties of flows and cuts

Residual graph

Augmenting path





MaxFlow MinCut Thm

Ford Fulkerson alg

Maximum matching in Bip graphs

Disjoint paths problem

# FF example



Max Flow and Min Cut

Properties of flows and cuts

Residual graph

Augmenting path

MaxFlow MinCut Thm

Ford Fulkerson alg

Maximum matching in Bip graphs

Disjoint paths problem

# FF example

Max Flow and  
Min Cut

Properties of  
flows and cuts

Residual  
graph


Augmenting  
path

MaxFlow  
MinCut Thm


Ford  
Fulkerson alg

Maximum  
matching in  
Bip graphs

Disjoint paths  
problem



Flow with max value



$\{s\}, \{a, b, t\}$  is a min  $(s, t)$ -cut

# Correctness of Ford-Fulkerson

Max Flow and  
Min Cut

Properties of  
flows and cuts

Residual  
graph

Augmenting  
path

MaxFlow  
MinCut Thm

**Ford  
Fulkerson alg**

Maximum  
matching in  
Bip graphs

Disjoint paths  
problem

Consequence of the Max-flow min-cut theorem.

## Theorem

*The flow returned by Ford-Fulkerson is the max-flow.*

# Networks with integer capacities

## Lemma (*Integrality invariant*)

Let  $\mathcal{N} = (V, E, c, s, t)$  where  $c : E \rightarrow \mathbb{Z}^+$ . At every iteration of the Ford-Fulkerson algorithm, the flow values  $f(e)$  are integers.

Proof: (induction)

- The statement is true for the initial flow (all zeroes).
- Inductive Hypothesis: The statement is true after  $j$  iterations.
- At iteration  $j + 1$ : As all residual capacities in  $G_f$  are integers, then bottleneck  $(P, f) \in \mathbb{Z}$ , for the augmenting path found in iteration  $j + 1$ .
- Thus the augmented flow values are integers. □

Max Flow and  
Min Cut

Properties of  
flows and cuts

Residual  
graph

Augmenting  
path

MaxFlow  
MinCut Thm

Ford  
Fulkerson alg

Maximum  
matching in  
Bip graphs

Disjoint paths  
problem

# Networks with integer capacities

Max Flow and  
Min Cut

Properties of  
flows and cuts

Residual  
graph

Augmenting  
path

MaxFlow  
MinCut Thm

Ford  
Fulkerson alg

Maximum  
matching in  
Bip graphs

Disjoint paths  
problem

## Theorem (**Integrality theorem**)

Let  $\mathcal{N} = (V, E, c, s, t)$  where  $c : E \rightarrow \mathbb{Z}^+$ . There exists a max-flow  $f^*$  such that  $f^*(e)$  is an integer, for any  $e \in E$ .

Proof:

Since the algorithm terminates, the theorem follows from the integrality invariant lemma. □



# Networks with integer capacities: FF running time

Max Flow and  
Min Cut

Properties of  
flows and cuts

Residual  
graph

Augmenting  
path

MaxFlow  
MinCut Thm

Ford  
Fulkerson alg

Maximum  
matching in  
Bip graphs

Disjoint paths  
problem

## Lemma

*Let  $C$  be the min cut capacity (=max. flow value),  
Ford-Fulkerson terminates after finding at most  $C$  augmenting  
paths.*

**Proof:** The value of the flow increases by  $\geq 1$  after each  
augmentation. □

# Networks with integer capacities: FF running time

Max Flow and  
Min Cut

Properties of  
flows and cuts

Residual  
graph

Augmenting  
path

MaxFlow  
MinCut Thm

Ford  
Fulkerson alg


Maximum  
matching in  
Bip graphs

Disjoint paths  
problem

- The number of iterations is  $\leq C$ . At each iteration:
- Constructing  $G_f$ , with  $E(G_f) \leq 2m$ , takes  $O(m)$  time.
- $O(n + m)$  time to find an augmenting path, or deciding that it does not exist.
- Total running time is  $O(C(n + m)) = O(Cm)$
- Is that polynomic? No, only pseudopolynomic

# Networks with integer capacities: FF running time


The number of iterations of Ford-Fulkerson could be  $\Theta(C)$



Ford-Fulkerson can alternate between the two long paths, and require  $2M$  iterations. Taking  $M = 10^{10}$ , FF on a graph with 4 vertices can take time  $2 \cdot 10^{10}$ .

# Networks with integer capacities: FF running time

The number of iterations of Ford-Fulkerson could be  $\Theta(C)$



Ford-Fulkerson can alternate between the two long paths, and require  $2M$  iterations. Taking  $M = 10^{10}$ , FF on a graph with 4 vertices can take time  $2 \cdot 10^{10}$ .

- 1 Max Flow and Min Cut
- 2 Properties of flows and cuts
- 3 Residual graph
- 4 Augmenting path
- 5 MaxFlow MinCut Thm
- 6 Ford Fulkerson alg
- 7 Maximum matching in Bip graphs**

Max Flow and  
Min Cut

Properties of  
flows and cuts

Residual  
graph

Augmenting  
path

MaxFlow  
MinCut Thm

Ford  
Fulkerson alg

**Maximum  
matching in  
Bip graphs**

Disjoint paths  
problem

# MAXIMUM MATCHING problem

Given an undirected graph  $G = (V, E)$  a subset of edges  $M \subseteq E$  is a **matching** if each node appears at most in one edge in  $M$  (a node may not appear at all).

**MAXIMUM MATCHING** problem:

Given a graph  $G$ , find a matching with maximum cardinality.

Max Flow and  
Min Cut

Properties of  
flows and cuts

Residual  
graph

Augmenting  
path

MaxFlow  
MinCut Thm

Ford  
Fulkerson alg

**Maximum  
matching in  
Bip graphs**


Disjoint paths  
problem

# MAXIMUM MATCHING problem

Given an undirected graph  $G = (V, E)$  a subset of edges  $M \subseteq E$  is a **matching** if each node appears at most in one edge in  $M$  (a node may not appear at all).

**MAXIMUM MATCHING** problem:

Given a graph  $G$ , find a matching with maximum cardinality.



Max Flow and  
Min Cut

Properties of  
flows and cuts

Residual  
graph

Augmenting  
path

MaxFlow  
MinCut Thm

Ford  
Fulkerson alg

Maximum  
matching in  
Bip graphs

Disjoint paths  
problem

# Maximum matching in bipartite graphs

A graph  $G = (V, E)$  is **bipartite** if there is a partition of  $V$  in  $L$  and  $R$ , ( $L \cup R = V$  and  $L \cap R = \emptyset$ ), such that every  $e \in E$  connects a vertex in  $L$  with a vertex in  $R$ .

Max Flow and  
Min Cut

Properties of  
flows and cuts

Residual  
graph

Augmenting  
path

MaxFlow  
MinCut Thm

Ford  
Fulkerson alg

**Maximum  
matching in  
Bip graphs**


Disjoint paths  
problem



# Maximum matching in bipartite graphs

A graph  $G = (V, E)$  is **bipartite** if there is a partition of  $V$  in  $L$  and  $R$ , ( $L \cup R = V$  and  $L \cap R = \emptyset$ ), such that every  $e \in E$  connects a vertex in  $L$  with a vertex in  $R$ .

We want to solve the **MAXIMUM MATCHING** problem on **bipartite graphs**



Max Flow and  
Min Cut

Properties of  
flows and cuts

Residual  
graph

Augmenting  
path

MaxFlow  
MinCut Thm

Ford  
Fulkerson alg


**Maximum  
matching in  
Bip graphs**

Disjoint paths  
problem

# Maximum matching in bipartite graphs

A graph  $G = (V, E)$  is **bipartite** if there is a partition of  $V$  in  $L$  and  $R$ , ( $L \cup R = V$  and  $L \cap R = \emptyset$ ), such that every  $e \in E$  connects a vertex in  $L$  with a vertex in  $R$ .

We want to solve the **MAXIMUM MATCHING** problem on **bipartite graphs**



Max Flow and  
Min Cut

Properties of  
flows and cuts

Residual  
graph

Augmenting  
path

MaxFlow  
MinCut Thm

Ford  
Fulkerson alg


Maximum  
matching in  
Bip graphs

Disjoint paths  
problem

# Maximum matching in bipartite graphs

A graph  $G = (V, E)$  is **bipartite** if there is a partition of  $V$  in  $L$  and  $R$ , ( $L \cup R = V$  and  $L \cap R = \emptyset$ ), such that every  $e \in E$  connects a vertex in  $L$  with a vertex in  $R$ .

We want to solve the **MAXIMUM MATCHING** problem on **bipartite graphs**



Max Flow and  
Min Cut

Properties of  
flows and cuts

Residual  
graph

Augmenting  
path

MaxFlow  
MinCut Thm

Ford  
Fulkerson alg



Maximum  
matching in  
Bip graphs

Disjoint paths  
problem

# MAXIMUM MATCHING: Network formulation

From  $G = (L \cup R, E)$  construct  $\mathcal{N} = (\hat{V}, \hat{E}, c, s, t)$ :

- Add vertices  $s$  and  $t$ :  $\hat{V} = L \cup R \cup \{s, t\}$ .
- Add directed edges  $s \rightarrow L$  with capacity 1. Add directed edges  $R \rightarrow t$  with capacity 1.
- Direct the edges  $E$  from  $L$  to  $R$ , and give them capacity  $\infty$ .
- $\hat{E} = \{s \rightarrow L\} \cup E \cup \{R \rightarrow t\}$ .



Max Flow and  
Min Cut

Properties of  
flows and cuts

Residual  
graph

Augmenting  
path

MaxFlow  
MinCut Thm

Ford  
Fulkerson alg

Maximum  
matching in  
Bip graphs

Disjoint paths  
problem

# Maximum matching algorithm: Analysis


## Theorem

*Max flow in  $\mathcal{N}$  = Max bipartite matching in  $G$ .*

### Proof Matching as flows

Let  $M$  be a matching in  $G$  with  $k$ -edges, consider the flow  $f$  that sends 1 unit along each one of the  $k$  paths,  $s \rightarrow u \rightarrow v \rightarrow t$ , for  $(u, v) \in M$ .

As  $M$  is a matching all these paths are disjoint, so  $f$  is a flow and has value  $k$ .



Max Flow and  
Min Cut

Properties of  
flows and cuts

Residual  
graph

Augmenting  
path

MaxFlow  
MinCut Thm

Ford  
Fulkerson alg

Maximum  
matching in  
Bip graphs

Disjoint paths  
problem

# Maximum matching algorithm: Analysis


## Theorem

*Max flow in  $\mathcal{N}$  = Max bipartite matching in  $G$ .*

## Proof Matching as flows

Let  $M$  be a matching in  $G$  with  $k$ -edges, consider the flow  $f$  that sends 1 unit along each one of the  $k$  paths,  $s \rightarrow u \rightarrow v \rightarrow t$ , for  $(u, v) \in M$ .


As  $M$  is a matching all these paths are disjoint, so  $f$  is a flow and has value  $k$ .



# Maximum matching algorithm: Analysis

## Flows as matchings

- Consider an integral flow  $f$  in  $\hat{G}$ . Therefore, for any edge  $e$ , the flow is either 0 or 1.
- Consider the cut  $C = (\{s\} \cup L, R \cup \{t\})$  in  $\hat{G}$ .
- Let  $M$  be the set of edges in the cut  $C$  with flow=1, then  $|M| = |f|$ .
- Each node in  $L$  is in at most one  $e \in M$  and every node in  $R$  is in at most one head of an  $e \in F$
- Therefore,  $M$  is a matching in  $G$  with  $|M| \leq |f|$



Max Flow and  
Min Cut

Properties of  
flows and cuts

Residual  
graph

Augmenting  
path

MaxFlow  
MinCut Thm

Ford  
Fulkerson alg



Maximum  
matching in  
Bip graphs

Disjoint paths  
problem

# Maximum matching algorithm: Analysis

## Flows as matchings

- Consider an integral flow  $f$  in  $\hat{G}$ . Therefore, for any edge  $e$ , the flow is either 0 or 1.
- Consider the cut  $C = (\{s\} \cup L, R \cup \{t\})$  in  $\hat{G}$ .
- Let  $M$  be the set of edges in the cut  $C$  with flow=1, then  $|M| = |f|$ .
- Each node in  $L$  is in at most one  $e \in M$  and every node in  $R$  is in at most one head of an  $e \in F$
- Therefore,  $M$  is a matching in  $G$  with  $|M| \leq |f|$





# Maximum matching algorithm: Analysis

As  $\mathcal{N}$  has integer capacities there is an integral maximum flow  $f^*$ , the associated matching is a maximum matching.  $\square$

Max Flow and  
Min Cut

Properties of  
flows and cuts

Residual  
graph

Augmenting  
path

MaxFlow  
MinCut Thm

Ford  
Fulkerson alg

**Maximum  
matching in  
Bip graphs**

Disjoint paths  
problem

# Maximum matching algorithm: Analysis

What is the cost of the algorithm?

Max Flow and  
Min Cut

Properties of  
flows and cuts

Residual  
graph

Augmenting  
path

MaxFlow  
MinCut Thm

Ford  
Fulkerson alg

**Maximum  
matching in  
Bip graphs**

Disjoint paths  
problem

# Maximum matching algorithm: Analysis

## What is the cost of the algorithm?

- The bipartite graph, has  $n$  vertices and  $m$  edges. The capacities are integers. We need an integral solution.
- The algorithm: (1) constructs  $\mathcal{N}$ , (2) runs FF on  $\mathcal{N}$  to obtain a maxflow  $f$ , (3) from  $f$  obtain a maximum matching  $M$ .

Max Flow and  
Min Cut

Properties of  
flows and cuts

Residual  
graph

Augmenting  
path

MaxFlow  
MinCut Thm

Ford  
Fulkerson alg

**Maximum  
matching in  
Bip graphs**

Disjoint paths  
problem

# Maximum matching algorithm: Analysis

## What is the cost of the algorithm?

- The bipartite graph, has  $n$  vertices and  $m$  edges. The capacities are integers. We need an integral solution.
- The algorithm: (1) constructs  $\mathcal{N}$ , (2) runs FF on  $\mathcal{N}$  to obtain a maxflow  $f$ , (3) from  $f$  obtain a maximum matching  $M$ .
- $\mathcal{N}$  has  $n + 2$  vertices and  $m + 2n$  edge, (1) takes  $O(n + m)$

Max Flow and  
Min Cut

Properties of  
flows and cuts

Residual  
graph

Augmenting  
path

MaxFlow  
MinCut Thm

Ford  
Fulkerson alg

Maximum  
matching in  
Bip graphs

Disjoint paths  
problem

# Maximum matching algorithm: Analysis

## What is the cost of the algorithm?

- The bipartite graph, has  $n$  vertices and  $m$  edges. The capacities are integers. We need an integral solution.
- The algorithm: (1) constructs  $\mathcal{N}$ , (2) runs FF on  $\mathcal{N}$  to obtain a maxflow  $f$ , (3) from  $f$  obtain a maximum matching  $M$ .
- $\mathcal{N}$  has  $n + 2$  vertices and  $m + 2n$  edge, (1) takes  $O(n + m)$
- The maximum value of a flow in  $\mathcal{N}$  is at most  $n$ , (2) takes time  $O(|f|(n + m)) = O(n(n + m))$

Max Flow and  
Min Cut

Properties of  
flows and cuts

Residual  
graph

Augmenting  
path

MaxFlow  
MinCut Thm

Ford  
Fulkerson alg

Maximum  
matching in  
Bip graphs

Disjoint paths  
problem

# Maximum matching algorithm: Analysis

## What is the cost of the algorithm?

- The bipartite graph, has  $n$  vertices and  $m$  edges. The capacities are integers. We need an integral solution.
- The algorithm: (1) constructs  $\mathcal{N}$ , (2) runs FF on  $\mathcal{N}$  to obtain a maxflow  $f$ , (3) from  $f$  obtain a maximum matching  $M$ .
- $\mathcal{N}$  has  $n + 2$  vertices and  $m + 2n$  edge, (1) takes  $O(n + m)$
- The maximum value of a flow in  $\mathcal{N}$  is at most  $n$ , (2) takes time  $O(|f|(n + m)) = O(n(n + m))$
- (3) can be done in time  $O(n + m)$ .

Max Flow and  
Min Cut

Properties of  
flows and cuts

Residual  
graph

Augmenting  
path

MaxFlow  
MinCut Thm

Ford  
Fulkerson alg

Maximum  
matching in  
Bip graphs

Disjoint paths  
problem

# Maximum matching algorithm: Analysis

## What is the cost of the algorithm?

- The bipartite graph, has  $n$  vertices and  $m$  edges. The capacities are integers. We need an integral solution.
- The algorithm: (1) constructs  $\mathcal{N}$ , (2) runs FF on  $\mathcal{N}$  to obtain a maxflow  $f$ , (3) from  $f$  obtain a maximum matching  $M$ .
- $\mathcal{N}$  has  $n + 2$  vertices and  $m + 2n$  edge, (1) takes  $O(n + m)$
- The maximum value of a flow in  $\mathcal{N}$  is at most  $n$ , (2) takes time  $O(|f|(n + m)) = O(n(n + m))$
- (3) can be done in time  $O(n + m)$ .

So, the cost is  $O(n(n + m))$ .

Max Flow and  
Min Cut

Properties of  
flows and cuts

Residual  
graph

Augmenting  
path

MaxFlow  
MinCut Thm

Ford  
Fulkerson alg

Maximum  
matching in  
Bip graphs

Disjoint paths  
problem

- 1 Max Flow and Min Cut
- 2 Properties of flows and cuts
- 3 Residual graph
- 4 Augmenting path
- 5 MaxFlow MinCut Thm
- 6 Ford Fulkerson alg
- 7 Maximum matching in Bip graphs

Max Flow and  
Min Cut

Properties of  
flows and cuts

Residual  
graph

Augmenting  
path

MaxFlow  
MinCut Thm

Ford  
Fulkerson alg

Maximum  
matching in  
Bip graphs


Disjoint paths  
problem



# DISJOINT PATH problem

Given a digraph  $G = (V, E)$  and two vertices  $s, t \in V$ , a set of paths is **edge-disjoint** if their edges are disjoint (although they might share some vertex)

**DISJOINT PATH problem:** Given a digraph  $G = (V, E)$  and two vertices  $s, t \in V$ , find a set of  $s \rightsquigarrow t$  edge-disjoint paths of maximum cardinality



Max Flow and  
Min Cut

Properties of  
flows and cuts

Residual  
graph

Augmenting  
path

MaxFlow  
MinCut Thm

Ford  
Fulkerson alg


Maximum  
matching in  
Bip graphs

Disjoint paths  
problem

# DISJOINT PATH: Max flow formulation

Thinking in terms of flow a path from  $s$  to  $t$  can be seen as a way of transporting a unit of flow.

We construct a network  $\mathcal{N}$  assigning unit capacity to every edge.



Max Flow and  
Min Cut

Properties of  
flows and cuts

Residual  
graph

Augmenting  
path

MaxFlow  
MinCut Thm

Ford  
Fulkerson alg

Maximum  
matching in  
Bip graphs


Disjoint paths  
problem

# DISJOINT PATH: Max flow formulation

Thinking in terms of flow a path from  $s$  to  $t$  can be seen as a way of transporting a unit of flow.

We construct a network  $\mathcal{N}$  assigning unit capacity to every edge

We solve MaxFlow for  $\mathcal{N}$ .



Max Flow and  
Min Cut

Properties of  
flows and cuts

Residual  
graph

Augmenting  
path

MaxFlow  
MinCut Thm

Ford  
Fulkerson alg

Maximum  
matching in  
Bip graphs

Disjoint paths  
problem

# DISJOINT PATH: Max flow formulation

Max Flow and  
Min Cut

Properties of  
flows and cuts

Residual  
graph

Augmenting  
path

MaxFlow  
MinCut Thm

Ford  
Fulkerson alg

Maximum  
matching in  
Bip graphs

Disjoint paths  
problem

Thinking in terms of flow a path from  $s$  to  $t$  can be seen as a way of transporting a unit of flow.

We construct a network  $\mathcal{N}$  assigning unit capacity to every edge

# DISJOINT PATH: Max flow formulation

Max Flow and  
Min Cut

Properties of  
flows and cuts

Residual  
graph

Augmenting  
path

MaxFlow  
MinCut Thm

Ford  
Fulkerson alg

Maximum  
matching in  
Bip graphs

Disjoint paths  
problem

Thinking in terms of flow a path from  $s$  to  $t$  can be seen as a way of transporting a unit of flow.

We construct a network  $\mathcal{N}$  assigning unit capacity to every edge

## Theorem

*The max number of edge disjoint paths  $s \rightsquigarrow t$  is equal to the max flow value*

# DISJOINT PATH: Proof of the Theorem

## Proof.

Number of disjoint paths  $\leq$  max flow

If we have  $k$  edge-disjoint paths  $s \rightsquigarrow t$  in  $G$  then making  $f(e) = 1$  for each  $e$  in a path, we get a flow with  $|f| = k$

Max Flow and  
Min Cut

Properties of  
flows and cuts

Residual  
graph

Augmenting  
path

MaxFlow  
MinCut Thm

Ford  
Fulkerson alg

Maximum  
matching in  
Bip graphs

Disjoint paths  
problem

# DISJOINT PATH: Proof of the Theorem

Number of disjoint paths  $\geq$  max flow

- If the max flow value is  $k$ , there exists a 0-1 flow  $f^*$  with value  $k$ .
- Consider the graph  $G^* = (V, E')$  where  $E'$  is formed by all edges  $e$  with  $f(e) = 1$ .
- We repeatedly compute a  $s \rightsquigarrow t$  simple path in  $G^*$ , and remove its edges from  $G^*$ .
- Each time that we remove a path, the value of the flow in the network is reduced by one, so we can apply the process  $k$  times.
- None of the paths share an edge, so we get  $k$  disjoint paths. □

Max Flow and  
Min Cut

Properties of  
flows and cuts

Residual  
graph

Augmenting  
path

MaxFlow  
MinCut Thm

Ford  
Fulkerson alg

Maximum  
matching in  
Bip graphs

Disjoint paths  
problem

# DISJOINT PATH: Proof of the Theorem

Number of disjoint paths  $\geq$  max flow

- If the max flow value is  $k$ , there exists a 0-1 flow  $f^*$  with value  $k$ .
- Consider the graph  $G^* = (V, E')$  where  $E'$  is formed by all edges  $e$  with  $f(e) = 1$ .
- We repeatedly compute a  $s \rightsquigarrow t$  simple path in  $G^*$ , and remove its edges from  $G^*$ .
- Each time that we remove a path, the value of the flow in the network is reduced by one, so we can apply the process  $k$  times.
- None of the paths share an edge, so we get  $k$  disjoint paths. □

End Proof

Max Flow and  
Min Cut

Properties of  
flows and cuts

Residual  
graph

Augmenting  
path

MaxFlow  
MinCut Thm

Ford  
Fulkerson alg

Maximum  
matching in  
Bip graphs

Disjoint paths  
problem



# DISJOINT PATH: Max flow + path extraction algorithm

## Algorithm

- 1 Construct the network  $\mathcal{N}$  assigning unit capacity to every edge
- 2 Solve MaxFlow for  $\mathcal{N}$
- 3 Extract the set of disjoint paths on the graph restricted to edges with flow  $> 0$

Max Flow and  
Min Cut

Properties of  
flows and cuts

Residual  
graph

Augmenting  
path

MaxFlow  
MinCut Thm

Ford  
Fulkerson alg


Maximum  
matching in  
Bip graphs

Disjoint paths  
problem

# DISJOINT PATH: Max flow + path extraction algorithm

## Algorithm

- 1 Construct the network  $\mathcal{N}$  assigning unit capacity to every edge
- 2 Solve MaxFlow for  $\mathcal{N}$
- 3 Extract the set of disjoint paths on the graph restricted to edges with flow  $> 0$



Max Flow and  
Min Cut

Properties of  
flows and cuts

Residual  
graph

Augmenting  
path

MaxFlow  
MinCut Thm

Ford  
Fulkerson alg


Maximum  
matching in  
Bip graphs

Disjoint paths  
problem

# DISJOINT PATH: Max flow + path extraction algorithm

## Algorithm

- 1 Construct the network  $\mathcal{N}$  assigning unit capacity to every edge
- 2 Solve MaxFlow for  $\mathcal{N}$
- 3 Extract the set of disjoint paths on the graph restricted to edges with flow  $> 0$



Max Flow and  
Min Cut

Properties of  
flows and cuts

Residual  
graph

Augmenting  
path

MaxFlow  
MinCut Thm

Ford  
Fulkerson alg

Maximum  
matching in  
Bip graphs

Disjoint paths  
problem

# Disjoint paths algorithm: Analysis

What is the cost of the algorithm?

Max Flow and  
Min Cut

Properties of  
flows and cuts

Residual  
graph

Augmenting  
path

MaxFlow  
MinCut Thm

Ford  
Fulkerson alg

Maximum  
matching in  
Bip graphs

**Disjoint paths  
problem**

# Disjoint paths algorithm: Analysis

## What is the cost of the algorithm?

- The graph, has  $n$  vertices and  $m$  edges. The capacities are integers. We need an integral solution.
- The algorithm: (1) constructs  $\mathcal{N}$ , (2) runs FF on  $\mathcal{N}$  to obtain a max flow  $f$ , (3) from  $f$  obtains  $|f|$  edge disjoint paths.

Max Flow and  
Min Cut

Properties of  
flows and cuts

Residual  
graph

Augmenting  
path

MaxFlow  
MinCut Thm

Ford  
Fulkerson alg

Maximum  
matching in  
Bip graphs

Disjoint paths  
problem

# Disjoint paths algorithm: Analysis

## What is the cost of the algorithm?

- The graph, has  $n$  vertices and  $m$  edges. The capacities are integers. We need an integral solution.
- The algorithm: (1) constructs  $\mathcal{N}$ , (2) runs FF on  $\mathcal{N}$  to obtain a max flow  $f$ , (3) from  $f$  obtains  $|f|$  edge disjoint paths.
- $\mathcal{N}$  has  $n$  vertices and  $m$  edges, (1) takes  $O(n + m)$

Max Flow and  
Min Cut

Properties of  
flows and cuts

Residual  
graph

Augmenting  
path

MaxFlow  
MinCut Thm

Ford  
Fulkerson alg

Maximum  
matching in  
Bip graphs

Disjoint paths  
problem

# Disjoint paths algorithm: Analysis

## What is the cost of the algorithm?

- The graph, has  $n$  vertices and  $m$  edges. The capacities are integers. We need an integral solution.
- The algorithm: (1) constructs  $\mathcal{N}$ , (2) runs FF on  $\mathcal{N}$  to obtain a max flow  $f$ , (3) from  $f$  obtains  $|f|$  edge disjoint paths.
- $\mathcal{N}$  has  $n$  vertices and  $m$  edges, (1) takes  $O(n + m)$
- The maximum value of a flow in  $\mathcal{N}$  is at most  $n$ , (2) takes time  $O(|f|(n + m)) = O(n(n + m))$

Max Flow and  
Min Cut

Properties of  
flows and cuts

Residual  
graph

Augmenting  
path

MaxFlow  
MinCut Thm

Ford  
Fulkerson alg

Maximum  
matching in  
Bip graphs

Disjoint paths  
problem

# Disjoint paths algorithm: Analysis

## What is the cost of the algorithm?

- The graph, has  $n$  vertices and  $m$  edges. The capacities are integers. We need an integral solution.
- The algorithm: (1) constructs  $\mathcal{N}$ , (2) runs FF on  $\mathcal{N}$  to obtain a max flow  $f$ , (3) from  $f$  obtains  $|f|$  edge disjoint paths.
- $\mathcal{N}$  has  $n$  vertices and  $m$  edges, (1) takes  $O(n + m)$
- The maximum value of a flow in  $\mathcal{N}$  is at most  $n$ , (2) takes time  $O(|f|(n + m)) = O(n(n + m))$
- (3) can be done in time  $O(n + m)$  per path, i.e.,  $O(|f|(n + m))$ .

Max Flow and  
Min Cut

Properties of  
flows and cuts

Residual  
graph

Augmenting  
path

MaxFlow  
MinCut Thm

Ford  
Fulkerson alg

Maximum  
matching in  
Bip graphs

Disjoint paths  
problem



# Disjoint paths algorithm: Analysis

## What is the cost of the algorithm?

- The graph, has  $n$  vertices and  $m$  edges. The capacities are integers. We need an integral solution.
- The algorithm: (1) constructs  $\mathcal{N}$ , (2) runs FF on  $\mathcal{N}$  to obtain a max flow  $f$ , (3) from  $f$  obtains  $|f|$  edge disjoint paths.
- $\mathcal{N}$  has  $n$  vertices and  $m$  edges, (1) takes  $O(n + m)$
- The maximum value of a flow in  $\mathcal{N}$  is at most  $n$ , (2) takes time  $O(|f|(n + m)) = O(n(n + m))$
- (3) can be done in time  $O(n + m)$  per path, i.e.,  $O(|f|(n + m))$ .

So the cost is  $O(n(n + m))$ .

Max Flow and  
Min Cut

Properties of  
flows and cuts

Residual  
graph

Augmenting  
path

MaxFlow  
MinCut Thm

Ford  
Fulkerson alg

Maximum  
matching in  
Bip graphs

Disjoint paths  
problem

# VERTEX DISJOINT PATHS

Can we do something similar to get the maximum number of vertex disjoint paths?

Max Flow and  
Min Cut

Properties of  
flows and cuts

Residual  
graph

Augmenting  
path

MaxFlow  
MinCut Thm

Ford  
Fulkerson alg

Maximum  
matching in  
Bip graphs

**Disjoint paths  
problem**

# The case of undirected graphs

If we have an undirected graph, with two distinguished nodes  $u, v$ , how would you apply the max flow formulation to solve the problem of finding the max number of disjoint paths between  $u$  and  $v$ ?

Max Flow and  
Min Cut

Properties of  
flows and cuts

Residual  
graph

Augmenting  
path

MaxFlow  
MinCut Thm

Ford  
Fulkerson alg

Maximum  
matching in  
Bip graphs

**Disjoint paths  
problem**

# The case of undirected graphs

Max Flow and  
Min Cut

Properties of  
flows and cuts

Residual  
graph

Augmenting  
path

MaxFlow  
MinCut Thm

Ford  
Fulkerson alg

Maximum  
matching in  
Bip graphs

**Disjoint paths  
problem**