Max-flow and min-cut problems

- **[Edmonds](#page-1-0)**
- [Generalized](#page-26-0)
- **[Circulations](#page-40-0)** [Lower bounds](#page-65-0)

[Edmonds](#page-1-0) Karp alg

[Generalized](#page-26-0)

[Circulations](#page-40-0) [Lower bounds](#page-65-0) [Joint rounding](#page-88-0)

[Min cost Max](#page-98-0) Flow

1 [Edmonds Karp alg](#page-1-0)

2 [Generalized assignment problems](#page-26-0)

3 [Circulations](#page-40-0)

4 [Min cost Max Flow](#page-98-0)

メロトメ 伊 トメ 君 トメ 君 トッ 重 $2Q$

Dinic and Edmonds-Karp algorithm

[Edmonds](#page-1-0) Karp alg

[Generalized](#page-26-0) assignment problems

[Circulations](#page-40-0) [Lower bounds](#page-65-0) [Joint rounding](#page-88-0)

[Min cost Max](#page-98-0) Flow

J.Edmonds, R. Karp: Theoretical improvements in algorithmic efficiency for network flow problems. Journal ACM 1972.

Yefim Dinic: Algorithm for solution of a problem of maximum flow in a network with power estimation. Doklady Ak.N. 1970

Choosing a good augmenting path can lead to a faster algorithm. Use BFS to find an augmenting paths in G_f .

∢ ロ ▶ . ∢ @ ▶ . ∢ ミ

Edmonds-Karp algorithm

[Edmonds](#page-1-0) Karp alg

[Generalized](#page-26-0) assignment problems

```
Circulations
Lower bounds
Joint rounding
```
[Min cost Max](#page-98-0) Flow

FF algorithm but using BFS: choose the augmenting path in G_f with the smallest length (number of edges).

```
Edmonds-Karp(G, c, s, t)For all e = (u, v) \in E let f(u, v) = 0G_f = Gwhile there is an s \rightarrow t path in G_fdo
  P = \mathsf{BFS}(G_f, s, t)f =Augment(f, P)
```
Compute G_f return f

 $choose: \rightarrow or$ The BFS in EK will

BFS paths on G_f

[Edmonds](#page-1-0) Karp alg

[Generalized](#page-26-0) assignment problems

[Circulations](#page-40-0) [Lower bounds](#page-65-0) [Joint rounding](#page-88-0)

[Min cost Max](#page-98-0) Flow

For $\mathcal{N} = (V, E, c, s, t)$ and a flow f in \mathcal{N} , assuming that G_f has an augmenting path, let f' be the next flow after executing one step of the EK algorithm.

メロト メタト メミト メミト

つくい

BFS paths on G_f

[Edmonds](#page-1-0) Karp alg

[Generalized](#page-26-0) assignment problems

[Circulations](#page-40-0) [Lower bounds](#page-65-0) [Joint rounding](#page-88-0)

[Min cost Max](#page-98-0) Flow

For $\mathcal{N} = (V, E, c, s, t)$ and a flow f in N, assuming that G_f has an augmenting path, let f' be the next flow after executing one step of the EK algorithm.

 \blacksquare The path from s to t in a BFS traversal starting at s, is a path $s \rightarrow t$ with minimum number of edges, i.e., a shortest length path.

K ロ ⊁ K 倒 ≯ K ミ ⊁ K ミ ≯

BFS paths on G_f

[Edmonds](#page-1-0) Karp alg

[Generalized](#page-26-0) assignment problems

[Circulations](#page-40-0) [Lower bounds](#page-65-0) [Joint rounding](#page-88-0)

[Min cost Max](#page-98-0) Flow

For $\mathcal{N} = (V, E, c, s, t)$ and a flow f in N, assuming that G_f has an augmenting path, let f' be the next flow after executing one step of the EK algorithm.

- \blacksquare The path from s to t in a BFS traversal starting at s, is a path $s \rightarrow t$ with minimum number of edges, i.e., a shortest length path.
- For \in V, let $\delta_f(s, v)$ denote length of a shortest length path from s to v in G_f .

∢ ロ ▶ 《 御 》 ∢ 重 》 ∢ 重 ▶ 。

[Edmonds](#page-1-0) Karp alg

[Generalized](#page-26-0)

[Circulations](#page-40-0) [Lower bounds](#page-65-0) [Joint rounding](#page-88-0)

[Min cost Max](#page-98-0) Flow

メロメメ 倒 メメ きょくきょう 目 299

[Edmonds](#page-1-0) Karp alg

[Generalized](#page-26-0) assignment problems

[Circulations](#page-40-0) [Lower bounds](#page-65-0) [Joint rounding](#page-88-0)

[Min cost Max](#page-98-0) Flow

How can we have $(u, v) \in E_{f'}$ but $(u, v) \notin E_f$? (u, v) is a forward edge saturated in f and not in f' . (u, v) is a backward edge in G_f and $f(v, u) = 0$

 $2Q$

メロメ メ団 メメミメメミメ

[Edmonds](#page-1-0) Karp alg

[Generalized](#page-26-0) assignment problems

[Circulations](#page-40-0) [Lower bounds](#page-65-0) [Joint rounding](#page-88-0)

[Min cost Max](#page-98-0) Flow

How can we have $(u, v) \in E_{f'}$ but $(u, v) \notin E_f$?

 (u, v) is a forward edge saturated in f and not in f' .

 (u, v) is a backward edge in G_f and $f(v, u) = 0$

In any of the two cases, the augmentation must have modified the flow from v to u, so (u, v) must form part of the augmenting path.

∢ロ ▶ ∢ 御 ▶ ∢ 后 ▶ ∢ 后 ▶

[Edmonds](#page-1-0) Karp alg

[Generalized](#page-26-0) assignment problems

[Circulations](#page-40-0) [Lower bounds](#page-65-0) [Joint rounding](#page-88-0)

[Min cost Max](#page-98-0) Flow

Lemma

If the EK-algorithm runs on $\mathcal{N} = (V, E, c, s, t)$, for all vertices $v \neq s$, $\delta_f(s, v)$ increases monotonically with each flow augmentation.

Proof. By contradiction.

Let f be the first flow such that, for some $u \neq s$,

 $\delta_{f'}(s, u) < \delta_f(s, u).$

メロメ メ団 メメミメメミメ

[Edmonds](#page-1-0) Karp alg

[Generalized](#page-26-0) assignment problems

[Circulations](#page-40-0) [Lower bounds](#page-65-0) [Joint rounding](#page-88-0)

[Min cost Max](#page-98-0) Flow

Proof (cont)

Let v be the vertex with the minimum $\delta_{f'}(s,v)$ whose distance was decreased.

Let $P: s \leadsto u \rightarrow v$ be a shortest length path from s to v in $G_{f'}$

Then, $\delta_{f'}(s,v) = \delta_{f'}(s,u) + 1$ and $\delta_{f'}(s,u) \geq \delta_f(s,u)$.

If $(u, v) \in E_f$, $\delta_f({\pmb s},\mathsf{v}) \leq \delta_f({\pmb s},\mathsf{u}) + 1 \leq \delta_{f'}({\pmb s},\mathsf{u}) + 1 = \delta_{f'}({\pmb s},\mathsf{v})$ ■ So, $(u, v) \notin E_f$

[Edmonds](#page-1-0) Karp alg

[Generalized](#page-26-0) assignment problems

[Circulations](#page-40-0) [Lower bounds](#page-65-0) [Joint rounding](#page-88-0)

[Min cost Max](#page-98-0) Flow

Proof (cont)

How can we have ?

 $(u,v) \in E_{f'}$ but $(u,v) \notin E_f$

If so, (v, u) appears in the augmenting path.

 $2Q$

メロメ メ団 メメミメメミメ

[Edmonds](#page-1-0) Karp alg

[Generalized](#page-26-0) assignment problems

[Circulations](#page-40-0) [Lower bounds](#page-65-0) [Joint rounding](#page-88-0)

[Min cost Max](#page-98-0) Flow

Proof (cont)

How can we have ?

 $(u,v) \in E_{f'}$ but $(u,v) \notin E_f$

If so, (v, u) appears in the augmenting path.

Then, the shortest length path from s to u in G_f **has** (v, u) as it last edge.

 $\delta_f(\pmb{s},\pmb{\nu}) \leq \delta_f(\pmb{s},\pmb{\nu}) - 1 \leq \delta_{f'}(\pmb{s},\pmb{\nu}) - 1 = \delta_{f'}(\pmb{s},\pmb{\nu}) - 1 - 1$

メロメ メ御 メメ ミメメ ミメ

[Edmonds](#page-1-0) Karp alg

[Generalized](#page-26-0) assignment problems

[Circulations](#page-40-0) [Lower bounds](#page-65-0) [Joint rounding](#page-88-0)

[Min cost Max](#page-98-0) Flow

Proof (cont)

How can we have ?

 $(u,v) \in E_{f'}$ but $(u,v) \notin E_f$

If so, (v, u) appears in the augmenting path.

Then, the shortest length path from s to u in G_f **has** (v, u) as it last edge.

$$
\delta_f(s,v) \leq \delta_f(s,u)-1 \leq \delta_{f'}(s,u)-1 = \delta_{f'}(s,v)-1-1
$$

メロメ メ団 メメミメメミメ

へのへ

which contradicts $\delta_{f'}(s, v) < \delta_f(u, v)$.

[Edmonds](#page-1-0) Karp alg

[Generalized](#page-26-0)

[Circulations](#page-40-0) [Lower bounds](#page-65-0) [Joint rounding](#page-88-0)

[Min cost Max](#page-98-0) Flow

メロメメ 倒 メメ きょくきょう 目 299

[Edmonds](#page-1-0) Karp alg

[Generalized](#page-26-0) assignment problems

[Circulations](#page-40-0) [Lower bounds](#page-65-0) [Joint rounding](#page-88-0)

[Min cost Max](#page-98-0) Flow

Let P be an augmenting path in G_f . $(u, v) \in P$ is critical if $b(P) = c_f(u, v)$.

メロメ メ御 メメ ミメメ ヨメ

 $2Q$

€

[Edmonds](#page-1-0) Karp alg

[Generalized](#page-26-0) assignment problems

[Circulations](#page-40-0) [Lower bounds](#page-65-0) [Joint rounding](#page-88-0)

[Min cost Max](#page-98-0) Flow

Let P be an augmenting path in G_f .

 $(u, v) \in P$ is critical if $b(P) = c_f(u, v)$.

Critical edges do not appear in $G_{f'}$.

$$
(u, v) \text{ forward}, f'(u, v) = c(u, v)
$$

 $2Q$

メロメ メ団 メメミメメミメ

 (u, v) backward, $f'(v, u) = 0$

Lemma

In the EK algorithm, each one of the edges can become critical at most $|V|/2$ times.

[Circulations](#page-40-0) [Lower bounds](#page-65-0) [Joint rounding](#page-88-0)

[Edmonds](#page-1-0) Karp alg [Generalized](#page-26-0) assignment problems

[Min cost Max](#page-98-0) Flow

Proof:

Let $(u, v) \in E$, when (u, v) is critical for the first time, $\delta_f(s, v) = \delta_f(s, u) + 1$

メロト メタト メミト メミト

つくい

Lemma

In the EK algorithm, each one of the edges can become critical at most $|V|/2$ times.

Proof:

- Let $(u, v) \in E$, when (u, v) is critical for the first time, $\delta_f(s, v) = \delta_f(s, u) + 1$
- After this step (u, v) disappears from the residual graph until after the flow in (u, v) changes.

∢ロ ▶ ∢ 御 ▶ ∢ 后 ▶ ∢ 后 ▶

へのへ

[Edmonds](#page-1-0) Karp alg

[Generalized](#page-26-0) assignment problems

[Circulations](#page-40-0) [Lower bounds](#page-65-0) [Joint rounding](#page-88-0)

Lemma

In the EK algorithm, each one of the edges can become critical at most $|V|/2$ times.

Proof:

- Let $(u, v) \in E$, when (u, v) is critical for the first time, $\delta_f(s, v) = \delta_f(s, u) + 1$
- After this step (u, v) disappears from the residual graph until after the flow in (u, v) changes.
- At this point, (v, u) forms part of the augmenting path in $G_{f'}$, and $\delta_{f'}(s, u) = \delta_{f'}(s, v) + 1$,

 $\delta_{f'}(\mathsf{s},\mathsf{u}) = \delta_{f'}(\mathsf{s},\mathsf{v}) + 1 \geq \delta_f(\mathsf{s},\mathsf{v}) + 1 \geq \delta_f(\mathsf{s},\mathsf{u}) + 2$

メロメ メ御 メメ ヨメメ ヨメー

へのへ

[Edmonds](#page-1-0) Karp alg

[Generalized](#page-26-0) assignment problems

[Circulations](#page-40-0) [Lower bounds](#page-65-0) [Joint rounding](#page-88-0)

Lemma

In the EK algorithm, each one of the edges can become critical at most $|V|/2$ times.

Proof:

- Let $(u, v) \in E$, when (u, v) is critical for the first time, $\delta_f(s, v) = \delta_f(s, u) + 1$
- After this step (u, v) disappears from the residual graph until after the flow in (u, v) changes.
- At this point, (v, u) forms part of the augmenting path in $G_{f'}$, and $\delta_{f'}(s, u) = \delta_{f'}(s, v) + 1$,

$$
\delta_{f'}(s,u)=\delta_{f'}(s,v)+1\geq \delta_f(s,v)+1\geq \delta_f(s,u)+2
$$

So, the distance has increased by [a](#page-20-0)t [le](#page-22-0)[a](#page-17-0)[s](#page-18-0)[t](#page-21-0) 2 [.](#page-0-0)

[Edmonds](#page-1-0) Karp alg

[Generalized](#page-26-0) assignment problems

[Circulations](#page-40-0) [Lower bounds](#page-65-0) [Joint rounding](#page-88-0)

Complexity of Edmonds-Karp algorithm

[Edmonds](#page-1-0) Karp alg

[Generalized](#page-26-0) assignment problems

[Circulations](#page-40-0) [Lower bounds](#page-65-0) [Joint rounding](#page-88-0)

[Min cost Max](#page-98-0) Flow

Theorem

The EK algorithms runs in $O(mn(n + m))$ steps. Therefore it is a polynomial time algorithm.

メロメ メ御 メメ ヨメメ ヨメー

重

 $2Q$

Complexity of Edmonds-Karp algorithm

[Edmonds](#page-1-0) Karp alg

[Generalized](#page-26-0) assignment problems

[Circulations](#page-40-0) [Lower bounds](#page-65-0) [Joint rounding](#page-88-0)

[Min cost Max](#page-98-0) Flow

Theorem

The EK algorithms runs in $O(mn(n + m))$ steps. Therefore it is a polynomial time algorithm.

Proof:

- Need time $O(m + n)$ to find the augmenting path using BFS.
- By the previous Lemma, there are $O(mn)$ augmentations. □

Finding a min-cut

[Edmonds](#page-1-0) Karp alg

[Generalized](#page-26-0) assignment problems

[Circulations](#page-40-0) [Lower bounds](#page-65-0) [Joint rounding](#page-88-0)

[Min cost Max](#page-98-0) Flow

- Given (G, s, t, c) to find a min-cut:
	- **1** Compute the max-flow f^* in G .
	- <mark>2</mark> Obtain *G_f*∗.
	- **3** Find the set $S = \{v \in V | s \leadsto v\}$ in $G_{f^*}.$
	- 4 Output the cut

 $(S, V - \{S\}) = \{(v, u) | v \in S \text{ and } u \in V - \{S\}\}\$ in G.

The running time is the same than the algorithm to find the max-flow.

メロト メタト メミト メミト

The max-flow problems: History

- Ford-Fulkerson (1956) $O(mC)$, where C is the max flow val.
- Dinic (1970) (blocking flow) $O(n^2m)$
- Edmond-Karp (1972) (shortest augmenting path) $O(nm^2)$
- Karzanov (1974), $O(n^2m)$ Goldberg-Tarjant (1986) (push re-label preflow $+$ dynamic trees) $O(nm\lg(n^2/m))$ (uses parallel implementation)
- King-Rao-Tarjan (1998) $O(nm \log_{m/n \lg n} n)$.
- J. Orlin (2013) $O(nm)$ (clever follow up to KRT-98)
- Chen, Kyng, Liu, Peng, Gutenberg, Sachdeva (2022) $O(m^{1+o(1)})$ (polynomially bounded integral capacities) You can read [Quanta Magazine](https://www.quantamagazine.org/researchers-achieve-absurdly-fast-algorithm-for-network-flow-20220608/) article.

[Edmonds](#page-1-0) Karp alg

[Generalized](#page-26-0) assignment problems

[Circulations](#page-40-0) [Lower bounds](#page-65-0) [Joint rounding](#page-88-0)

[Edmonds](#page-1-0) Karp alg

[Generalized](#page-26-0) assignment problems

[Circulations](#page-40-0) [Lower bounds](#page-65-0) [Joint rounding](#page-88-0)

[Min cost Max](#page-98-0) Flow

1 [Edmonds Karp alg](#page-1-0)

2 [Generalized assignment problems](#page-26-0)

3 [Circulations](#page-40-0)

4 [Min cost Max Flow](#page-98-0)

メロトメ 伊 トメ 君 トメ 君 トッ 重 $2Q$

Final's Scheduling

We have as input:

- \blacksquare n courses, each one with a final. Each exam must be given in one room. Each course c_i has $E[i]$ students.
- r rooms. Each r_i has a capacity $S[i]$,
- \blacksquare τ time slots. For each room and time slot, we only can schedule one final.
- \blacksquare p professors to watch exams. Each exam needs one professor in each class and time. Each professor has its own restrictions of availability and no professor can oversee more than 6 finals. For each p_ℓ and τ_k define a Boolean variable $A[k,\ell]=\mathcal{T}$ if p_ℓ is available at $\tau_k.$

Design an efficient algorithm that correctly schedules a room, a time slot and a professor to every final, or report that not such schedule is possible.メロト メタト メミト メミト つくい

[Edmonds](#page-1-0) Karp alg

[Generalized](#page-26-0) assignment problems

[Circulations](#page-40-0) [Lower bounds](#page-65-0) [Joint rounding](#page-88-0)

Construction of the network

[Edmonds](#page-1-0) Karp alg

[Generalized](#page-26-0) assignment problems

[Circulations](#page-40-0) [Lower bounds](#page-65-0) [Joint rounding](#page-88-0)

[Min cost Max](#page-98-0) Flow

Construct the network N with vertices $\{s, t, \{c_i\}, \{r_i\}, \{\tau_k\}, \{p_\ell\}\}\$. Edges and capacities:

- (s, c_i) with capacity 1 (each course has one final)
- (c_i,r_j) , if $E[i]\leq S[j]$, with capacity ∞ (can go to a room in which it fits)
- $\forall j,k,$ $(r_j,\tau_k),$ with capacity 1 (one final per room and time slot).
- (τ_k, p_ℓ) , if $A[k, \ell] = T$, capacity 1 (p can watch one final, if p is available at τ_k). (ρ_ℓ,t) , capacity 6 (each ρ can watch ≤ 6 finals) Notice that neither rooms nor time slots have individual restrictions.

Final's Scheduling: Flow Network

[Edmonds](#page-1-0) Karp alg

[Generalized](#page-26-0) assignment problems

[Circulations](#page-40-0) [Lower bounds](#page-65-0)

[Min cost Max](#page-98-0)

courses rooms

slots professors

メロメメ 倒 トメ 君 トメ 君 トー \equiv 299

Final's Scheduling

[Edmonds](#page-1-0) Karp alg

[Generalized](#page-26-0) assignment problems

[Circulations](#page-40-0) [Lower bounds](#page-65-0) [Joint rounding](#page-88-0)

- Notice the input size to the problem is
	- $N = n + r + \tau + p + 2$. and size of the network is $O(N)$ vertices and $O(N^2)$ edges, why?
- Every path $s \leadsto t$ is an assignment of room-time-professor to a final, and any assignment room-time-professor to a final can be represented by a path $s \leadsto t$.
- Every integral flow identifies a collection of $|f|$ (s, t) -paths leading to a valid assignment for $|f|$ finals and viceversa.

Final's Scheduling

[Edmonds](#page-1-0) Karp alg

[Generalized](#page-26-0) assignment problems

[Circulations](#page-40-0) [Lower bounds](#page-65-0) [Joint rounding](#page-88-0)

[Min cost Max](#page-98-0) Flow

- To maximize the number of finals to be given, we compute the max-flow f^* from s to t .
- If $|f^*| = n$, then we can schedule all finals, otherwise we can not.
- To recover the assignment we have to consider the edges with positive flow and extract assignment from the n (s, t) -paths
- Complexity:
	- To construct $\mathcal N$, we need $O(N^2)$.
	- As $|f^*| \le n$ integral, we can use Ford-Fulkerson to compute f^* , with cost $O(nN^2)$.

K ロ ⊁ K 倒 ≯ K ミ ⊁ K ミ ≯

∽≏ດ

- The second part requires $O(N^2)$ time.
- So, the cost of the algorithm is $O(nN^2) = O(n(n + r + \tau + p)^2).$

[Edmonds](#page-1-0) Karp alg

[Generalized](#page-26-0) assignment problems

[Circulations](#page-40-0) [Lower bounds](#page-65-0) [Joint rounding](#page-88-0)

[Min cost Max](#page-98-0) Flow

Gonsider a generalized assignment problem \mathcal{GP} where, we have as input d finite sets X_1, \ldots, X_d , each representing a different set of resources.

メロメ メ御 メメ ミメメ ミメ

つくい

- **[Edmonds](#page-1-0)** Karp alg
- [Generalized](#page-26-0) assignment problems
- **[Circulations](#page-40-0)** [Lower bounds](#page-65-0) [Joint rounding](#page-88-0)
- [Min cost Max](#page-98-0) Flow
- Gonsider a generalized assignment problem \mathcal{GP} where, we have as input d finite sets X_1, \ldots, X_d , each representing a different set of resources.
- \blacksquare Our goal is to chose the "largest" number of d-tuples, each d -tuple containing exactly one element from each $X_i,$ subject to the constrains:

K ロ ⊁ K 倒 ≯ K ミ ⊁ K ミ ≯

- **[Edmonds](#page-1-0)** Karp alg
- [Generalized](#page-26-0) assignment problems
- **[Circulations](#page-40-0)**
- [Lower bounds](#page-65-0) [Joint rounding](#page-88-0)
- [Min cost Max](#page-98-0) Flow
- Gonsider a generalized assignment problem \mathcal{GP} where, we have as input d finite sets X_1, \ldots, X_d , each representing a different set of resources.
- \blacksquare Our goal is to chose the "largest" number of d-tuples, each d -tuple containing exactly one element from each $X_i,$ subject to the constrains:
	- **■** For each $i \in [d]$, each $x \in X_i$ can appears in at most $c(x)$ selected tuples.
	- For each $i \in [d]$, any two $x \in X_i$ and $y \in X_{i+1}$ can appear in at most $c(x, y)$ selected tuples.

メロメ メ御 メメ ミメメ ミメ

∽≏ດ

The values for $c(x)$ and $c(x, y)$ are either in \mathbb{Z}^+ or ∞ .

- **[Edmonds](#page-1-0)** Karp alg
- [Generalized](#page-26-0) assignment problems
- **[Circulations](#page-40-0)** [Lower bounds](#page-65-0)
- [Joint rounding](#page-88-0)
- [Min cost Max](#page-98-0) Flow
- Gonsider a generalized assignment problem \mathcal{GP} where, we have as input d finite sets X_1, \ldots, X_d , each representing a different set of resources.
- \blacksquare Our goal is to chose the "largest" number of d-tuples, each d -tuple containing exactly one element from each $X_i,$ subject to the constrains:
	- **■** For each $i \in [d]$, each $x \in X_i$ can appears in at most $c(x)$ selected tuples.
	- For each $i \in [d]$, any two $x \in X_i$ and $y \in X_{i+1}$ can appear in at most $c(x, y)$ selected tuples.
	- The values for $c(x)$ and $c(x, y)$ are either in \mathbb{Z}^+ or ∞ .
	- Notice that only pairs of objects between adjacent X_i and X_{i+1} are constrained.
Applications: Generic reduction to Max-Flow

[Edmonds](#page-1-0) Karp alg

[Generalized](#page-26-0) assignment problems

[Circulations](#page-40-0) [Lower bounds](#page-65-0) [Joint rounding](#page-88-0)

[Min cost Max](#page-98-0) Flow

Make the reduction from \mathcal{GP} to the following network \mathcal{N} :

- V contains a vertex x , for each element x in each X_i , and a copy x' , for each element $x \in X_i$ for $1 \leq i < d$.
- \blacksquare We add vertex s and vertex t.
- Add an edge $s \to x$ for each $x \in X_1$ and add an edge $y \to t$ for every $y \in X_d$. Give capacities $c(s, x) = c(x)$ and $c(y, t) = c(y)$.
- Add an edge $x' \to y$ for every pair $x \in X_i$ and $y \in X_{i+1}$. Give a capacity $c(x, y)$. Omit the edges with capacity 0.
- For every $x \in X_i$ for $1 \leq i < d$, add an edge $x \to x'$ with $c(x, x') = c(x)$.

Every path $s \leadsto t$ in $\mathcal N$ identifies a feasible d-tuple, conversely every d-[t](#page-35-0)uple determines a a path $s \rightsquigarrow t$ [.](#page-37-0) へのへ

Flow Network: The reduction

[Edmonds](#page-1-0) Karp alg

[Generalized](#page-26-0) assignment problems

[Circulations](#page-40-0) [Lower bounds](#page-65-0)

[Min cost Max](#page-98-0) Flow

メロメメ 御 メメ ミメメ ミメ È 299

Flow Network: The reduction

[Edmonds](#page-1-0) Karp alg

[Generalized](#page-26-0) assignment problems

[Circulations](#page-40-0) [Lower bounds](#page-65-0) [Joint rounding](#page-88-0)

- To solve \mathcal{GP} , we construct \mathcal{N} , and then we find an integer maximum flow f^* .
- In the subgraph formed by edges with $f^*(e) > 0$, we find a (s, t) path P (a d-tuple), decrease in 1 the flow in each edge of P , remove edges with 0 flow.

Flow Network: The reduction

[Edmonds](#page-1-0) Karp alg

[Generalized](#page-26-0) assignment problems

[Circulations](#page-40-0) [Lower bounds](#page-65-0) [Joint rounding](#page-88-0)

- To solve \mathcal{GP} , we construct \mathcal{N} , and then we find an integer maximum flow f^* .
- In the subgraph formed by edges with $f^*(e) > 0$, we find a (s, t) path P (a d-tuple), decrease in 1 the flow in each edge of P , remove edges with 0 flow.
- Repeat $|f^*|$ times. In this way we obtain a set of d-tuples with maximum size verifying all [the](#page-38-0) [re](#page-40-0)[s](#page-40-0)[tr](#page-37-0)[i](#page-39-0)[c](#page-40-0)[ti](#page-25-0)[o](#page-26-0)[n](#page-39-0)s[.](#page-25-0) Ω

[Edmonds](#page-1-0) Karp alg

[Generalized](#page-26-0)

[Circulations](#page-40-0)

[Lower bounds](#page-65-0) [Joint rounding](#page-88-0)

[Min cost Max](#page-98-0) Flow

1 [Edmonds Karp alg](#page-1-0)

2 [Generalized assignment problems](#page-26-0)

3 [Circulations](#page-40-0)

4 [Min cost Max Flow](#page-98-0)

K ロ ▶ K 御 ▶ K 君 ▶ K 君 ▶ . 重 $2Q$

Circulation with demands

[Edmonds](#page-1-0) Karp alg

[Generalized](#page-26-0) assignment problems

[Circulations](#page-40-0) [Demands](#page-41-0)

[Lower bounds](#page-65-0) [Joint rounding](#page-88-0)

[Min cost Max](#page-98-0) Flow

■ We introduce another flow problem, to deal with supply and demand inside a network.

 $2Q$

∢ロ ▶ ∢ 御 ▶ ∢ 后 ▶ ∢ 后 ▶

Instead of having a pair source/sink the new setting $\mathcal{L}_{\mathcal{A}}$ consider a producer/consumer scenario.

Circulation with demands

[Edmonds](#page-1-0) Karp alg

[Generalized](#page-26-0) assignment problems

[Circulations](#page-40-0) [Demands](#page-41-0)

[Lower bounds](#page-65-0) [Joint rounding](#page-88-0)

- We introduce another flow problem, to deal with supply and demand inside a network.
- **Instead of having a pair source/sink the new setting** consider a producer/consumer scenario.
- Some nodes are able to produce a certain amount of flow.
- Some nodes are willing to consume flow.
- \blacksquare The question is whether it is possible to route "all" the produced flow to the consumers. When possible the flow assignment is called a circulation

Network with demands

[Edmonds](#page-1-0) Karp alg

[Generalized](#page-26-0) assignment problems

[Circulations](#page-40-0) [Demands](#page-41-0)

[Lower bounds](#page-65-0) [Joint rounding](#page-88-0)

[Min cost Max](#page-98-0) Flow

A network with demands N is a tuple (V, E, c, d) where c assigns a positive capacity to each edge, and d is a function associating a demand $d(v)$, to $v \in V$.

 $2Q$ ∢ロ ▶ ∢ 御 ▶ ∢ 后 ▶ ∢ 后 ▶

Network with demands

A network with demands N is a tuple (V, E, c, d) where c assigns a positive capacity to each edge, and d is a function associating a demand $d(v)$, to $v \in V$.

[Circulations](#page-40-0) [Demands](#page-41-0) [Lower bounds](#page-65-0) [Joint rounding](#page-88-0) [Min cost Max](#page-98-0) Flow

[Edmonds](#page-1-0) Karp alg [Generalized](#page-26-0) assignment problems

- When $d(v) > 0$, v can receive $d(v)$ units of flow more than it sends, v is a sink,.
- If $d(v) < 0$, v can send $d(v)$ units of flow more than it receives, v is a source.
- If $d(v) = 0$, v is neither a source or a sink.

Network with demands

A network with demands N is a tuple (V, E, c, d) where c assigns a positive capacity to each edge, and d is a function associating a demand $d(v)$, to $v \in V$.

[Circulations](#page-40-0) [Demands](#page-41-0) [Lower bounds](#page-65-0) [Joint rounding](#page-88-0)

[Edmonds](#page-1-0) Karp alg [Generalized](#page-26-0) assignment problems

- When $d(v) > 0$, v can receive $d(v)$ units of flow more than it sends, v is a sink,.
- If $d(v) < 0$, v can send $d(v)$ units of flow more than it receives, v is a source.
- If $d(v) = 0$, v is neither a source or a sink.
- Define S to be the set of sources and T the set of sinks.

Network with demands: circulation

[Edmonds](#page-1-0) Karp alg

[Generalized](#page-26-0) assignment problems

[Circulations](#page-40-0) [Demands](#page-41-0)

[Lower bounds](#page-65-0) [Joint rounding](#page-88-0)

[Min cost Max](#page-98-0) Flow

Given a network $\mathcal{N} = (V, E, c, d)$, a circulation is a flow assignment $f: E \to \mathbb{R}^+$ s.t.

1 capacity: For each $e \in E$, $0 \leq f(e) \leq c(e)$,

2 conservation: For each $v \in V$,

へのへ

メロメ メ団 メメミメメミメ

Network with demands: circulation

[Edmonds](#page-1-0) Karp alg

[Generalized](#page-26-0) assignment problems

[Circulations](#page-40-0) [Demands](#page-41-0)

[Lower bounds](#page-65-0) [Joint rounding](#page-88-0)

[Min cost Max](#page-98-0) Flow

Given a network $\mathcal{N} = (V, E, c, d)$, a circulation is a flow assignment $f: E \to \mathbb{R}^+$ s.t.

1 capacity: For each $e \in E$, $0 \leq f(e) \leq c(e)$,

2 conservation: For each $v \in V$,

Take into account that a circulation might not exist.

Network with demands: circulation problem

Circulation problem: Given $\mathcal{N} = (V, E, c, d)$ with $c > 0$, obtain a circulation provided it does exists.

K ロ ▶ K 御 ▶ K 舌

 $2Q$

 $\left\{ \begin{array}{c} 1 \end{array} \right.$

[Edmonds](#page-1-0) Karp alg

[Generalized](#page-26-0) assignment problems

[Circulations](#page-40-0) [Demands](#page-41-0)

[Lower bounds](#page-65-0) [Joint rounding](#page-88-0)

If f is a circulation for $\mathcal{N} = (V, E, c, d)$,

$$
\sum_{v \in V} d(v) = \sum_{v \in V} \left(\underbrace{\sum_{(u,v) \in E} f(u,v)}_{\text{edges to } v} - \underbrace{\sum_{(v,z) \in E} f(v,z)}_{\text{edges out of } v} \right)
$$

.

重

 $2Q$

メロメ メ御 メメ ミメメ ミメー

[Edmonds](#page-1-0) Karp alg

[Generalized](#page-26-0) assignment

[Circulations](#page-40-0) [Demands](#page-41-0)

[Lower bounds](#page-65-0) [Joint rounding](#page-88-0)

If f is a circulation for $\mathcal{N} = (V, E, c, d)$,

$$
\sum_{v \in V} d(v) = \sum_{v \in V} \left(\underbrace{\sum_{(u,v) \in E} f(u,v)}_{\text{edges to } v} - \underbrace{\sum_{(v,z) \in E} f(v,z)}_{\text{edges out of } v} \right)
$$

.

へのへ

For $e = (u, v) \in E$, $f(e)$ appears in the sum of edges to v and in the sum of edges out of u . Both terms cancel!

 \leftarrow \cap \rightarrow \leftarrow \leftarrow \cap \rightarrow

[Edmonds](#page-1-0) Karp alg

[Generalized](#page-26-0) assignment problems

[Circulations](#page-40-0) [Demands](#page-41-0)

[Lower bounds](#page-65-0) [Joint rounding](#page-88-0)

If f is a circulation for $\mathcal{N} = (V, E, c, d)$,

$$
\sum_{v \in V} d(v) = \sum_{v \in V} \left(\underbrace{\sum_{(u,v) \in E} f(u,v)}_{\text{edges to } v} - \underbrace{\sum_{(v,z) \in E} f(v,z)}_{\text{edges out of } v} \right)
$$

.

へのへ

[Joint rounding](#page-88-0) [Min cost Max](#page-98-0) Flow

[Edmonds](#page-1-0) Karp alg [Generalized](#page-26-0) assignment problems **[Circulations](#page-40-0)** [Demands](#page-41-0) [Lower bounds](#page-65-0)

> For $e = (u, v) \in E$, $f(e)$ appears in the sum of edges to v and in the sum of edges out of u . Both terms cancel!

> > 4 m \rightarrow 4 m \rightarrow \rightarrow \pm

Then, $\sum_{v \in V} d(v) = 0$.

メロメ メ御 メメ ミメメ ミメー

重

 $2Q$

[Edmonds](#page-1-0) Karp alg

[Generalized](#page-26-0) assignment

[Circulations](#page-40-0) [Demands](#page-41-0)

[Lower bounds](#page-65-0) [Joint rounding](#page-88-0)

[Min cost Max](#page-98-0) Flow

If there is a circulation, then $\sum_{v \in V} d(v) = 0$.

Recall that

$$
S = \{v \in V | d(v) < 0\} \text{ and } T = \{v \in V | d(v) > 0\}.
$$

Define
$$
D = -\sum_{v \in S} d(v) = \sum_{v \in T} d(v)
$$
.

[Edmonds](#page-1-0) Karp alg

[Generalized](#page-26-0) assignment problems

[Circulations](#page-40-0) [Demands](#page-41-0)

[Lower bounds](#page-65-0) [Joint rounding](#page-88-0)

[Min cost Max](#page-98-0) Flow

If there is a circulation, then $\sum_{v \in V} d(v) = 0$.

Recall that

$$
S = \{v \in V | d(v) < 0\} \text{ and } T = \{v \in V | d(v) > 0\}.
$$

Define $D = -\sum_{v \in S} d(v) = \sum_{v \in T} d(v)$.

D is the total amount of extra flow that has to be transported from the sources to the sinks.

へのへ

 $\left\{ \begin{array}{ccc} 1 & 0 & 0 \\ 0 & 1 & 0 \end{array} \right\}$

[Edmonds](#page-1-0) Karp alg

[Generalized](#page-26-0) assignment problems

[Circulations](#page-40-0) [Demands](#page-41-0)

[Lower bounds](#page-65-0) [Joint rounding](#page-88-0)

[Min cost Max](#page-98-0) Flow

From $\mathcal{N} = (V, E, c, d)$, define a flow network $\mathcal{N}'=(\mathcal{V}', \mathcal{E}', \mathcal{c}', \mathsf{s}, t)$:

 $V' = V \cup \{s, t\}$, we add a source s and a sink t.

For $v \in S$ ($d(v) < 0$), add (s, v) with capacity $-d(v)$.

イロメ イタメ オラメイラメ

へのへ

For $v \in \mathcal{T}$ (d(v) > 0), add (v, t) with capacity d(v).

Keep E and, for $e \in E$, $c'(e) = c(e)$.

[Edmonds](#page-1-0) Karp alg

[Generalized](#page-26-0) assignment problems

[Circulations](#page-40-0) [Demands](#page-41-0)

[Lower bounds](#page-65-0) [Joint rounding](#page-88-0)

[Min cost Max](#page-98-0) Flow

From $\mathcal{N} = (V, E, c, d)$, define a flow network $\mathcal{N}'=(\mathcal{V}', \mathcal{E}', \mathcal{c}', \mathsf{s}, t)$:

 $V' = V \cup \{s, t\}$, we add a source s and a sink t.

For $v \in S$ ($d(v) < 0$), add (s, v) with capacity $-d(v)$.

For $v \in \mathcal{T}$ (d(v) > 0), add (v, t) with capacity d(v).

Keep E and, for $e \in E$, $c'(e) = c(e)$.

1.- Every flow f' in \mathcal{N}' verifies $|f'| \leq D$

[Edmonds](#page-1-0) Karp alg

[Generalized](#page-26-0) assignment

[Circulations](#page-40-0) [Demands](#page-41-0)

[Lower bounds](#page-65-0) [Joint rounding](#page-88-0)

[Min cost Max](#page-98-0) Flow

メロメ メ御 メメ ヨメメ ヨメー 重 $2Q$

1.- Every flow f' in \mathcal{N}' verifies $|f'| \leq D$

The capacity $c'(\{s\}, V) = D$, by the capacity restriction on flows, $|f'| \leq D$.

メロメ メ御 メメ ミメメ ミメ

 $2Q$

[Edmonds](#page-1-0) Karp alg

[Generalized](#page-26-0) assignment problems

[Circulations](#page-40-0) [Demands](#page-41-0)

[Lower bounds](#page-65-0) [Joint rounding](#page-88-0)

2.- If there is a circulation f in $\mathcal N$, we have a max-flow f' in $\mathcal N'$ with $|f'| = D$.

メロト メタト メミト メミト

つくい

Extend f to a flow f', assigning $f'(s, v) = -d(v)$, for $v \in S$, and $f'(u,t) = d(u)$, for $u \in \mathcal{T}$.

[Edmonds](#page-1-0) Karp alg

[Generalized](#page-26-0) assignment problems

[Circulations](#page-40-0) [Demands](#page-41-0)

[Lower bounds](#page-65-0) [Joint rounding](#page-88-0)

2.- If there is a circulation f in $\mathcal N$, we have a max-flow f' in $\mathcal N'$ with $|f'| = D$.

Extend f to a flow f', assigning $f'(s, v) = -d(v)$, for $v \in S$, and $f'(u,t) = d(u)$, for $u \in \mathcal{T}$.

By the circulation condition, f' is a flow in \mathcal{N}' . Furthermore, $|f'|=D.$

[Joint rounding](#page-88-0) [Min cost Max](#page-98-0) Flow

[Edmonds](#page-1-0) Karp alg [Generalized](#page-26-0) assignment problems **[Circulations](#page-40-0)** [Demands](#page-41-0) [Lower bounds](#page-65-0)

4 n + 4 n +

 Ω

Analysis

[Edmonds](#page-1-0) Karp alg

[Generalized](#page-26-0) assignment problems

[Circulations](#page-40-0) [Demands](#page-41-0)

[Lower bounds](#page-65-0) [Joint rounding](#page-88-0)

[Min cost Max](#page-98-0) Flow

3.- If there is a flow f' in \mathcal{N}' with $|f'| = D$, $\mathcal N$ has a circulation

For $e \in E$, define $f(e) = f'(e)$.

- As $|f'| = D$, all edges $(s, v) \in E'$ and $(u, t) \in E'$ are saturated by f' .
- \blacksquare By flow conservation, f satisfies $d(v) =$ $\sum f(u, v) - \sum f(v, z)$. $(u,v) \in E$ $=$ $\frac{d}{dx}$ $\frac{d}{dx}$ $\frac{d}{dx}$ $(v,z) \in E$ ${edges out of } v$

So, f is a circulation for N .

From the previous discussion, we can conclude:

Theorem (Necessary and sufficient condition)

There is a circulation for $\mathcal{N} = (V, E, c, d)$ iff the maxflow in \mathcal{N}' has value D.

メロメ メ御 メメ ミメメ ミメー

 $2Q$

店

[Edmonds](#page-1-0) Karp alg

[Generalized](#page-26-0) assignment problems

[Circulations](#page-40-0) [Demands](#page-41-0)

[Lower bounds](#page-65-0) [Joint rounding](#page-88-0)

From the previous discussion, we can conclude:

Theorem (Necessary and sufficient condition)

There is a circulation for $\mathcal{N} = (V, E, c, d)$ iff the maxflow in N' has value D

Theorem (Circulation integrality theorem)

If all capacities and demands are integers, and there exists a circulation, then there exists an integer valued circulation.

Sketch Proof Max-flow formulation $+$ integrality theorem for max -flow \Box

[Edmonds](#page-1-0) Karp alg

[Generalized](#page-26-0) assignment problems

[Circulations](#page-40-0) [Demands](#page-41-0)

[Lower bounds](#page-65-0) [Joint rounding](#page-88-0)

[Edmonds](#page-1-0) Karp alg

[Generalized](#page-26-0) assignment problems

[Circulations](#page-40-0) [Demands](#page-41-0)

[Lower bounds](#page-65-0) [Joint rounding](#page-88-0)

[Min cost Max](#page-98-0) Flow

Theorem

There is a polynomial time algorithm to solve the circulation problem.

 Ω

メロメ メ団 メメミメメミメ

The cost of the algorithm is the same as the cost of the algorithm used for the MaxFlow computation.

[Edmonds](#page-1-0) Karp alg

[Generalized](#page-26-0) assignment problems

[Circulations](#page-40-0) [Demands](#page-41-0)

[Lower bounds](#page-65-0) [Joint rounding](#page-88-0)

[Min cost Max](#page-98-0) Flow

Theorem

There is a polynomial time algorithm to solve the circulation problem.

The cost of the algorithm is the same as the cost of the algorithm used for the MaxFlow computation.

Theorem

If all capacities and demands are integers, and there exists a circulation, then we can obtain an integer valued circulation in time O(Dm).

Networks with demands and lower bounds

Generalization of the previous problem: besides satisfy demands at nodes, we want to force the flow to use certain edges.

メロメ メ御 メメ ミメメ ミメ

 $2Q$

店

[Edmonds](#page-1-0) Karp alg

[Generalized](#page-26-0) assignment problems

[Circulations](#page-40-0) [Lower bounds](#page-65-0) [Joint rounding](#page-88-0)

Networks with demands and lower bounds

Generalization of the previous problem: besides satisfy demands at nodes, we want to force the flow to use certain edges.

[Edmonds](#page-1-0) Karp alg

assignment problems

[Circulations](#page-40-0) [Lower bounds](#page-65-0) [Joint rounding](#page-88-0)

[Min cost Max](#page-98-0) Flow

Introduce a new constrain $\ell(e)$ on each $e \in E$, indicating the min-value the flow must be on e.

メロメ メ団 メメミメメミメ

 Ω

Networks with demands and lower bounds

Generalization of the previous problem: besides satisfy demands at nodes, we want to force the flow to use certain edges.

Introduce a new constrain $\ell(e)$ on each $e \in E$, indicating the min-value the flow must be on e.

A network $\mathcal N$ with demands and lower bounds is a tuple (V, E, c, ℓ, d) with $c(e) \geq \ell(e) \geq 0$, for each $e \in E$,

∢ロ ▶ ∢ 御 ▶ ∢ 后 ▶ ∢ 后 ▶

へのへ

[Edmonds](#page-1-0) Karp alg

[Generalized](#page-26-0) assignment problems

[Circulations](#page-40-0) [Lower bounds](#page-65-0) [Joint rounding](#page-88-0)

Networks with demands and lower bounds: circulation

[Edmonds](#page-1-0) Karp alg

[Generalized](#page-26-0) assignment problems

[Circulations](#page-40-0) [Lower bounds](#page-65-0) [Joint rounding](#page-88-0)

[Min cost Max](#page-98-0) Flow

Given a network $\mathcal{N} = (V, E, c, \ell, d)$ a circulation as a flow assignment $f: E \to \mathbb{R}^+$ s.t.

1 capacity: For each $e \in E$, $\ell(e) < f(e) < c(e)$,

2 conservation: For each $v \in V$,

 $\sum f(u, v) - \sum f(v, z) = d(v).$ (u,v)∈E (v,z)∈E

へのへ

A circulation might not exist.

Circulations with demands and lower bounds problem

Circulation with demands and lower bounds problem: Given $\mathcal{N} = (V, E, c, \ell, d)$, obtain a circulation for \mathcal{N} , provided it does exists

[Generalized](#page-26-0) assignment problems

[Edmonds](#page-1-0) Karp alg

[Circulations](#page-40-0) [Lower bounds](#page-65-0) [Joint rounding](#page-88-0)

[Min cost Max](#page-98-0) Flow

∢ ロ ▶ ∢ 伊 ▶ ∢ 舌

 \sim メミト $2Q$

Circulations with demands and lower bounds problem

Circulation with demands and lower bounds problem: Given $\mathcal{N} = (V, E, c, \ell, d)$, obtain a circulation for \mathcal{N} , provided it does exists

We devise an algorithm to the problem by a reduction to a circulation with demands problem.

メロト メタト メミト メミト

 $2Q$

[Edmonds](#page-1-0) Karp alg

[Generalized](#page-26-0) assignment problems

[Circulations](#page-40-0) [Lower bounds](#page-65-0) [Joint rounding](#page-88-0)

Circulations with demands and lower bounds: the reduction

Let $\mathcal{N} = (V, E, c, \ell, d)$, construct a network $\mathcal{N}'=(\mathcal{V}, \mathcal{E}, \mathsf{c}', \mathsf{d}')$ with only demands as follows:

[Edmonds](#page-1-0) Karp alg

[Generalized](#page-26-0) assignment problems

[Circulations](#page-40-0) [Lower bounds](#page-65-0) [Joint rounding](#page-88-0)

[Min cost Max](#page-98-0) Flow

Initially set $c' = c$ and $d' = d$. For each $e = (u, v) \in E$, with $\ell(e) > 0$: $c'(e) = c(e) - \ell(e).$

Update the demands on both ends of e : $d'(u) = d(u) + \ell(e)$ and $d'(v) = d(v) - \ell(e)$

へのへ
1.- If f is a circulation in $\mathcal N$, $f'(e) = f(e) - \ell(e)$, for $e \in E$, is a circulation in \mathcal{N}^{\prime} .

メロト メタト メミト メミト

 $2Q$

€

[Edmonds](#page-1-0) Karp alg

[Generalized](#page-26-0) assignment problems

[Circulations](#page-40-0) [Lower bounds](#page-65-0) [Joint rounding](#page-88-0)

1.- If f is a circulation in $\mathcal N$, $f'(e) = f(e) - \ell(e)$, for $e \in E$, is a circulation in \mathcal{N}^{\prime} .

By construction of \mathcal{N}' , f' verifies the capacity constraint.

Besides, for (u, v) with $\ell(u, v) > 0$, the flow out of u and the flow in v is decreased by $\ell(u, v)$.

f is a circulation in $\mathcal N$ so, the flow imbalance of f' matches the demand d' at each node.

[Edmonds](#page-1-0) Karp alg

[Generalized](#page-26-0) assignment problems

[Circulations](#page-40-0) [Lower bounds](#page-65-0) [Joint rounding](#page-88-0)

2.- If f' is a circulation in $\mathcal{N}',\ f(e)=f'(e)+\ell(e),$ for $e\in E,$ is a circulation in N .

メロト メタト メミト メミト

 $2Q$

€

[Edmonds](#page-1-0) Karp alg

[Generalized](#page-26-0) assignment problems

[Circulations](#page-40-0) [Lower bounds](#page-65-0) [Joint rounding](#page-88-0)

2.- If f' is a circulation in $\mathcal{N}',\ f(e)=f'(e)+\ell(e),$ for $e\in E,$ is a circulation in N .

 f' verifies the capacity constraint $f'(e) \geq 0$, so $f'(e) \geq \ell(e).$

 f' is a circulation, the f' imbalance at u is $d'(u)$.

Therefore, for (u, v) with $\ell(u, v) > 0$, the increase of flow in (u, v) balances $\ell(u, v)$ units of flow out of u with $\ell(u, v)$ units of flow entering v. Thus the f imbalance at u is $d(u)$.

[Edmonds](#page-1-0) Karp alg

[Generalized](#page-26-0) assignment problems

[Circulations](#page-40-0) [Lower bounds](#page-65-0) [Joint rounding](#page-88-0)

Main result

[Generalized](#page-26-0) assignment problems

[Circulations](#page-40-0) [Lower bounds](#page-65-0) [Joint rounding](#page-88-0)

[Min cost Max](#page-98-0) Flow

Theorem

There exists a circulation in N iff there exists a circulation in \mathcal{N}' . Moreover, if all demands, capacities and lower bounds in N are integers, and N admits a circulation, there is a circulation in N that is integer-valued.

The integer-valued circulation part is a consequence of the integer-value circulation Theorem for f' in G' .

K ロ ⊁ K 倒 ≯ K ミ ⊁ K ミ ≯

へのへ

Circulation with demands and lower bounds: main results

Theorem

[Edmonds](#page-1-0) Karp alg

[Generalized](#page-26-0) assignment problems

[Circulations](#page-40-0) [Lower bounds](#page-65-0) [Joint rounding](#page-88-0)

[Min cost Max](#page-98-0) Flow

There is a polynomial time algorithm to solve the circulation with demands and lower bounds problem.

The cost of the algorithm is the same as the cost of the algorithm used for the circulation with demands computation.

∢ロ ▶ ∢ 御 ▶ ∢ 后 ▶ ∢ 后 ▶

へのへ

Circulation with demands and lower bounds: main results

Theorem

[Edmonds](#page-1-0) Karp alg

[Generalized](#page-26-0) assignment problems

[Circulations](#page-40-0) [Lower bounds](#page-65-0) [Joint rounding](#page-88-0)

[Min cost Max](#page-98-0) Flow

There is a polynomial time algorithm to solve the circulation with demands and lower bounds problem.

The cost of the algorithm is the same as the cost of the algorithm used for the circulation with demands computation.

Theorem

If all capacities, lower bounds, and demands are integers, and there exists a circulation, then we can obtain an integer valued circulation in time $O((D + L)m)$ where L is the sum of al lower bounds.

SURVEY DESIGN problem

Problem: Design a survey among customers of products (KT-7.8)

- **Customer i can only be** asked about a bought product and must receive a questionnaire for at least c_i such products, those values are determined as function of the purchased products.
- For each product i , we want to collect data from a minimum of p_i customers.
- \blacksquare The c and p values cannot exceed the number of

purchased, sold produ[c](#page-0-0)ts.

へのへ

 \leftarrow \leftarrow

[Edmonds](#page-1-0) Karp alg

[Generalized](#page-26-0) assignment problems

[Circulations](#page-40-0) [Survey design](#page-79-0) [Joint rounding](#page-88-0)

SURVEY DESIGN problem

[Edmonds](#page-1-0) Karp alg

[Generalized](#page-26-0) assignment problems

[Circulations](#page-40-0) [Lower bounds](#page-65-0) [Survey design](#page-79-0) [Joint rounding](#page-88-0)

[Min cost Max](#page-98-0) Flow

The input to the problem is:

A set C of n customers and a set P of m products.

- For each customer $i \in \mathcal{C}$, a list of purchased products and the two values $c_i \leq c'_i$.
- For each product $j \in P$, two values p_j and p'_j .

Alternatively,

■ The information about purchases can be represented as a bipartite graph $G = (C \cup P, E)$, where C is the set of customers and P is the set of products.

■ $(i, j) \in E$ means $i \in C$ has purchased product $j \in P$.

SURVEY DESIGN: Input

[Edmonds](#page-1-0) Karp alg

[Generalized](#page-26-0)

[Circulations](#page-40-0) [Lower bounds](#page-65-0) [Survey design](#page-79-0)

Customers
$$
C = \{a, b, c, d\}
$$

Products $P = \{1, 2, 3, 4, 5, 6\}$

Survey Design: Circulation with lower bounds formulation

[Edmonds](#page-1-0) Karp alg

[Generalized](#page-26-0) assignment problems

[Circulations](#page-40-0) [Lower bounds](#page-65-0) [Survey design](#page-79-0) [Joint rounding](#page-88-0)

[Min cost Max](#page-98-0) Flow

We construct a network $\mathcal{N}=(\mathcal{V}',E',c,\ell)$ from G as follows:

- Nodes: $V' = V \cup \{s, t\}$
- Edges: E' contains E and edges $s \to \{C\}$, $\{P\} \to t$, and $(t,s).$
- Capacities and lower bounds:
	- c(t, s) = ∞ and $\ell(t,s) = 0$
	- For $i \in \mathcal{C}$, $\ell(s, i) = c_i$ and $c(s, i) =$ the number of purchased products.
		- For $j \in P$, $\ell(j, t) = p_i$ and $c(j, t) =$ number of customers that purchased j.
	- For $(i, j) \in E$, $c(i, j) = 1$, and $\ell(i, j) = 0$.

Survey Design: Circulation with lower bounds formulation

[Edmonds](#page-1-0) Karp alg

[Generalized](#page-26-0) assignment problems

[Circulations](#page-40-0) [Lower bounds](#page-65-0) [Survey design](#page-79-0) [Joint rounding](#page-88-0)

[Min cost Max](#page-98-0) Flow

Customers $C = \{a, b, c, c\}$ Products $P = \{1, 2, 3, 4, 5, 6\}$

Survey Design: Circulation interpretation

[Edmonds](#page-1-0) Karp alg

[Generalized](#page-26-0) assignment problems

[Circulations](#page-40-0) [Lower bounds](#page-65-0) [Survey design](#page-79-0) [Joint rounding](#page-88-0)

[Min cost Max](#page-98-0) Flow

If f is a circulation in N :

- one unit of flow circulates
	- $s \rightarrow i \rightarrow j \rightarrow t \rightarrow s$.
- **f** $f(i, j) = 1$ means ask i about j ,
- **f** $(f(s, i) \#$ products to ask i for opinion,
- **f** $f(i,t) = #$ customers to be asked to review i ,
- **f** $f(t,s)$ is the total number of questionnaires.

Survey Design: Circulation vs solutions

- [Edmonds](#page-1-0) Karp alg
- [Generalized](#page-26-0) assignment problems
- **[Circulations](#page-40-0)** [Lower bounds](#page-65-0) [Survey design](#page-79-0) [Joint rounding](#page-88-0)
- [Min cost Max](#page-98-0) Flow

A solution

- Ask a about $1, 2$.
- Ask *b* about $1, 2, 4$.
- Ask c about 3, 6.
- Ask d about 3, 4, 5.

3 로

∢ ロ ▶ . ∢ @ ▶ . ∢ ミ

 $2Q$

Main result

Theorem N has a circulation iff there is a feasible way to design the survey.

Proof if there is a feasible way to design the survey:

- **i** if *i* is asked about *j* then $f(i, j) = 1$,
- **f** $f(s, i)$ = number questions asked to i ($\geq c_i$).
- **f** (f, t) = number of customers who were asked about j $(> d_i),$

$f(t,s) =$ total number of questions.

E easy to verify that f is a circulation in \mathcal{N}

If there is an integral circulation in \mathcal{N} :

if $f(i, j) = 1$ then i will be asked about j,

 \blacksquare the constrains will be satisfied by the capacity rule.

メロメ メ御 メメ ヨメメ ヨメー

 Ω

[Edmonds](#page-1-0) Karp alg

[Generalized](#page-26-0) assignment problems

[Circulations](#page-40-0) [Lower bounds](#page-65-0) [Survey design](#page-79-0) [Joint rounding](#page-88-0)

Cost of the algorithm

- [Edmonds](#page-1-0) Karp alg
- [Generalized](#page-26-0) assignment problems
- **[Circulations](#page-40-0)** [Lower bounds](#page-65-0) [Survey design](#page-79-0) [Joint rounding](#page-88-0)
- [Min cost Max](#page-98-0) Flow
- \blacksquare N has $N = n + m + 2$ vertices and $E = n + m + nm$ edges
- $L = \sum_{e} \ell(e) \le nm$.
- \blacksquare Obtain $\mathcal N$ and extract the information from the circulation has cost $O(nm)$.

K ロ ⊁ K 倒 ≯ K ミ ⊁ K ミ ≯

へのへ

- \blacksquare FF analysis, the cost of obtaining a circulaton $O(L(N + M)) = O(n^2m^2)$.
- EK analysis, the cost of obtaining a circulaton $O(NM(N + M)) = O((n + m)n^2m^2)$.
- The algorithm has cost $O(n^2m^2)$.

[Edmonds](#page-1-0) Karp alg

[Generalized](#page-26-0) assignment problems

[Circulations](#page-40-0) [Lower bounds](#page-65-0) [Joint rounding](#page-88-0)

[Min cost Max](#page-98-0) Flow

Consider a matrix $A = (a_{ij})$ with dimension $n \times n$, where each $a_{ij} \in \mathbb{R}^+ \cup \{0\}$ and furthermore the sum of each row/column is an integer value. We want to round, if possible, each value a_{ii} to $|a_{ii}|$ or $[a_{ii}]$ without changing the value of the sum per row/column.

∢ロ ▶ ∢ 御 ▶ ∢ 后 ▶ ∢ 后 ▶

へのへ

Consider a matrix $A = (a_{ij})$ with dimension $n \times n$, where each $a_{ij} \in \mathbb{R}^+ \cup \{0\}$ and furthermore the sum of each row/column is an integer value. We want to round, if possible, each value a_{ii} to $|a_{ii}|$ or $[a_{ii}]$ without changing the value of the sum per row/column.

$$
\begin{pmatrix}\n10.9 & 2.5 & 1.3 & 9.3 \\
3.8 & 9.2 & 2.2 & 11.8 \\
7.9 & 5.2 & 7.3 & 0.6 \\
3.4 & 13.1 & 1.2 & 6.3\n\end{pmatrix} F \rightarrow \begin{pmatrix}\n11 & 3 & 1 & 9 \\
4 & 9 & 2 & 12 \\
7 & 5 & 8 & 1 \\
4 & 13 & 2 & 6\n\end{pmatrix}
$$

Such a rounding is called a joint rounding.

[Edmonds](#page-1-0) Karp alg

[Generalized](#page-26-0) assignment problems

[Circulations](#page-40-0) [Lower bounds](#page-65-0) [Joint rounding](#page-88-0)

Note that:

 \blacksquare The elements in A that are integers cannot be modified.

メロメ メ御 メメ ミメメ ヨメ

 $2Q$

店

Let $r_i = \sum_{j=1}^n (a_{ij} - \lfloor a_{ij} \rfloor)$ and $c_j = \sum_{i=1}^n (a_{ij} - \lfloor a_{ij} \rfloor)$

[Edmonds](#page-1-0) Karp alg

[Generalized](#page-26-0) assignment problems

[Circulations](#page-40-0) [Lower bounds](#page-65-0) [Joint rounding](#page-88-0)

Note that:

- \blacksquare The elements in A that are integers cannot be modified.
- Let $r_i = \sum_{j=1}^n (a_{ij} \lfloor a_{ij} \rfloor)$ and $c_j = \sum_{i=1}^n (a_{ij} \lfloor a_{ij} \rfloor)$
- As the rows/columns of A add up to an integer, r_i and c_i are integer values.

メロメ メ団 メメミメメミメ

へのへ

Furthermore,
$$
\sum_i r_i = \sum_j c_j
$$
.

[Edmonds](#page-1-0) Karp alg

[Generalized](#page-26-0) assignment problems

[Circulations](#page-40-0) [Lower bounds](#page-65-0) [Joint rounding](#page-88-0)

Note that:

- \blacksquare The elements in A that are integers cannot be modified.
	- Let $r_i = \sum_{j=1}^n (a_{ij} \lfloor a_{ij} \rfloor)$ and $c_j = \sum_{i=1}^n (a_{ij} \lfloor a_{ij} \rfloor)$
- As the rows/columns of A add up to an integer, r_i and c_i are integer values.

Furthermore,
$$
\sum_i r_i = \sum_j c_j
$$
.

In order to solve the problem we perform a reduction from this problem to a circulation problem.

∢ロ ▶ ∢ 御 ▶ ∢ 后 ▶ ∢ 后 ▶

へのへ

[Edmonds](#page-1-0) Karp alg

[Generalized](#page-26-0) assignment problems

[Circulations](#page-40-0) [Lower bounds](#page-65-0) [Joint rounding](#page-88-0)

Note that:

- \blacksquare The elements in A that are integers cannot be modified.
	- Let $r_i = \sum_{j=1}^n (a_{ij} \lfloor a_{ij} \rfloor)$ and $c_j = \sum_{i=1}^n (a_{ij} \lfloor a_{ij} \rfloor)$
- As the rows/columns of A add up to an integer, r_i and c_i are integer values.

Furthermore,
$$
\sum_i r_i = \sum_j c_j
$$
.

[Joint rounding](#page-88-0) [Min cost Max](#page-98-0) Flow

[Edmonds](#page-1-0) Karp alg [Generalized](#page-26-0) assignment problems **[Circulations](#page-40-0)** [Lower bounds](#page-65-0)

> In order to solve the problem we perform a reduction from this problem to a circulation problem.

One unit of flow can be seen as indicating that a decimal part rounded up to 1.

Zero flow indicates that a decimal part is rounded down to 0.

[Generalized](#page-26-0) assignment problems

[Circulations](#page-40-0) [Lower bounds](#page-65-0) [Joint rounding](#page-88-0)

[Min cost Max](#page-98-0) Flow

Build the network with demands $\mathcal{N} = (V, E, c, d)$ where:

 $\textsf{Vertices:}\;\; V = \{ \textit{x}_i, \textit{y}_i | 1 \leq i \leq \textit{n} \}$. \textit{x} 's vertices represent rows and y vertices columns.

メロメ メ御 メメ ヨメメ ヨメー

へのへ

- **Edges:** $E = \{(x_i, y_j) | 1 \le i, j \le n \text{ i } a_{i,j} \notin \mathbb{Z}\}\$
- **Capacities:** $c(x_i, y_j) = 1$.
- Demands: $d(x_i) = -r_i$, $1 \leq i \leq n$, i $d(y_j) = c_j$, $1 \leq j \leq n$.

 ${\cal N}$ has $O(n)$ vertices and $O(n^2)$ edges.

[Generalized](#page-26-0) assignment problems

[Circulations](#page-40-0) [Lower bounds](#page-65-0) [Joint rounding](#page-88-0)

[Min cost Max](#page-98-0) Flow

If there is a joint rounding of A, $\mathcal N$ has a circulation with integer values in $\{0, 1\}$.

 \blacksquare Let B a joint rounding of A, we define a new matrix D where

$$
d_{ij} = \begin{cases} 1 & \text{if } b_{ij} > a_{ij} \\ 0 & \text{otherwise} \end{cases}
$$

As B is a joint rounding, $\sum_j d_{ij} = r_i$ i $\sum_i d_{ij} = c_i$. Therefore, the flow assignment $f(i, j) = d_{ij}$ is a circulation in $\mathcal N$

[Generalized](#page-26-0) problems

[Circulations](#page-40-0) [Lower bounds](#page-65-0) [Joint rounding](#page-88-0)

[Min cost Max](#page-98-0) Flow

If N has a circulation with integer values in $\{0,1\}$, there is a joint rounding of A.

Let f be a circulation in N .

 \blacksquare we define matrix B as

$$
b_{i,j} = \begin{cases} a_{ij} & \text{if } a_{ij} \in \mathbb{Z} \\ \begin{bmatrix} a_{i,j} \end{bmatrix} & \text{if } a_{ij} \notin \mathbb{Z} \text{ if } f(i,j) = 1 \\ \begin{bmatrix} a_{i,j} \end{bmatrix} & \text{otherwise} \end{cases}
$$

As f is a circulation, $\sum_j b_{ij} = \sum_j a_{ij}$ and $\sum_i b_{ij} = \sum_i a_{ij}.$ **Therefore, B** is a joint rounding of A.

[Generalized](#page-26-0) assignment problems

[Circulations](#page-40-0) [Lower bounds](#page-65-0) [Joint rounding](#page-88-0)

[Min cost Max](#page-98-0) Flow

The construction of $\mathcal N$ has cost $O(n^2)$.

Ford-Fulkerson algorithm requires $O(D|E|)$, where D is the sum of the positive demands, i.e., $D=\sum r_i=O(n^2).$ As $|E| = O(n^2)$, the total cost is $O(n^4)$.

メロメ メ団 メメミメメミメ

つくい

[Generalized](#page-26-0)

[Circulations](#page-40-0) [Lower bounds](#page-65-0) [Joint rounding](#page-88-0)

[Min cost Max](#page-98-0) Flow

1 [Edmonds Karp alg](#page-1-0)

2 [Generalized assignment problems](#page-26-0)

3 [Circulations](#page-40-0)

4 [Min cost Max Flow](#page-98-0)

メロトメ 伊 トメ 君 トメ 君 トッ 重 $2Q$

Flow Network with costs

[Edmonds](#page-1-0) Karp alg

[Generalized](#page-26-0) assignment problems

[Circulations](#page-40-0) [Lower bounds](#page-65-0) [Joint rounding](#page-88-0)

[Min cost Max](#page-98-0) Flow

A network $\mathcal{N} = (V, E, c, \$, s, t)$ is formed by

- **a** digraph $G = (V, E)$,
- a source vertex $s \in V$
- **a** sink vertex $t \in V$,
- and edge capacities $c: E \rightarrow \mathbb{R}^+$
- and unit flow cost $\texttt{\$}:\textit{E}\rightarrow\text{I\!R}^+$

A flow in a network

Given a network $\mathcal{N} = (V, E, c, s, t)$ A Flow is an assignment $f:E\to\mathbb{R}^+\cup\{0\}$ that follows the Kirchoff's laws:

 $\forall (u, v) \in E, 0 \leq f(u, v) \leq c(u, v),$

(Flow conservation) $\forall v \in V - \{s, t\}$, $\sum_{u\in V} f(u,v) = \sum_{z\in V} f(v,z)$

The value of a flow f is

 $|f| = \sum f(s, v) = f(s, V) = f(V, t).$ v∈V

つくい

∢ロ ▶ ∢ 御 ▶ ∢ 后 ▶ ∢ 后 ▶

The cost of a flow f is

$$
\mathfrak{F}(f)=\sum_{e\in E}\mathfrak{F}(e)f(e).
$$

[Edmonds](#page-1-0) Karp alg

[Generalized](#page-26-0) assignment problems

[Circulations](#page-40-0) [Lower bounds](#page-65-0) [Joint rounding](#page-88-0)

A flow in a network

Given a network $\mathcal{N} = (V, E, c, s, t)$ A Flow is an assignment $f:E\to\mathbb{R}^+\cup\{0\}$ that follows the Kirchoff's laws:

 $\forall (u, v) \in E, 0 \leq f(u, v) \leq c(u, v),$

(Flow conservation) $\forall v \in V - \{s, t\}$, $\sum_{u\in V} f(u,v) = \sum_{z\in V} f(v,z)$

The value of a flow f is

 $|f| = \sum f(s, v) = f(s, V) = f(V, t).$ v∈V

The cost of a flow f is

$$
\mathfrak{F}(f)=\sum_{e\in E}\mathfrak{F}(e)f(e).
$$

[Edmonds](#page-1-0) Karp alg

[Generalized](#page-26-0) assignment problems

[Circulations](#page-40-0) [Lower bounds](#page-65-0) [Joint rounding](#page-88-0)

INPUT: A network flow with costs $\mathcal{N} = (V, E, c, \text{\textsterling}, s, t,)$ QUESTION: Find a flow of maximum value on N having minimum cost.

メロメ メ御 メメ ミメメ ミメ

 $2Q$

€

[Edmonds](#page-1-0) Karp alg

[Generalized](#page-26-0) assignment problems

[Circulations](#page-40-0) [Lower bounds](#page-65-0) [Joint rounding](#page-88-0)

INPUT: A network flow with costs $\mathcal{N} = (V, E, c, \text{\textsterling}, s, t,)$ QUESTION: Find a flow of maximum value on N having minimum cost.

 $\left\{ \begin{array}{ccc} 1 & 0 & 0 \\ 0 & 1 & 0 \end{array} \right\}$

 $2Q$

Red edges have unit cost 0.5 and all others unit cost 0.1

[Edmonds](#page-1-0) Karp alg

[Generalized](#page-26-0) assignment problems

[Circulations](#page-40-0) [Lower bounds](#page-65-0) [Joint rounding](#page-88-0)

INPUT: A network flow with costs $\mathcal{N} = (V, E, c, \text{\textsterling}, s, t,)$ QUESTION: Find a flow of maximum value on N having minimum cost.

Red edges have unit cost 0.5 and all others unit cost 0.1

 $|f| = 7$ (it is maximum) $\$(f) = 1.9 + 2 = 3.9$

[Edmonds](#page-1-0) Karp alg

[Generalized](#page-26-0) assignment problems

[Circulations](#page-40-0) [Lower bounds](#page-65-0) [Joint rounding](#page-88-0)

INPUT: A network flow with costs $\mathcal{N} = (V, E, c, \text{\textsterling}, s, t,)$ QUESTION: Find a flow of maximum value on N having minimum cost.

 $\left\{ \begin{array}{ccc} 1 & 0 & 0 \\ 0 & 1 & 0 \end{array} \right\}$

 $2Q$

Red edges have unit cost 0.5 and all others unit cost 0.1

[Edmonds](#page-1-0) Karp alg

[Generalized](#page-26-0) assignment problems

[Circulations](#page-40-0) [Lower bounds](#page-65-0) [Joint rounding](#page-88-0)

INPUT: A network flow with costs $\mathcal{N} = (V, E, c, \text{\textsterling}, s, t,)$ QUESTION: Find a flow of maximum value on N having minimum cost.

Red edges have unit cost 0.5 and all others unit cost 0.1

 $|f| = 7$ (it is maximum) $\$(f) = 2.4 + 0.5 = 3.4$

メロト メタト メミト メミト

 $2Q$

[Edmonds](#page-1-0) Karp alg

[Generalized](#page-26-0) assignment problems

[Circulations](#page-40-0) [Lower bounds](#page-65-0) [Joint rounding](#page-88-0)

Flows and cycles in the residual graph

[Edmonds](#page-1-0) Karp alg

[Generalized](#page-26-0) assignment problems

[Circulations](#page-40-0) [Lower bounds](#page-65-0) [Joint rounding](#page-88-0)

[Min cost Max](#page-98-0) Flow

Given a network with costs $\mathcal{N} = (V, E, s, \mathsf{\$}, t, c)$ together with a flow f on it, the residual graph, $(G_f=(V,E_f,c_f,\$_f)$ is a weighted digraph on the same vertex set and with edge set:

- **i** if $c(u, v) f(u, v) > 0$, then $(u, v) \in E_f$ and $c_f(u, v) = c(u, v) - f(u, v) > 0$ and $\oint(u, v) = \oint(u, v)$ (forward edges)
- **■** if $f(u, v) > 0$, then $(v, u) \in E_f$ and $c_f(v, u) = f(u, v)$ and $\$(v, u) = -\(u, v) (backward edges).

Let C be a (simple) cycle in G_f , the bottleneck, $b(C)$, is the minimum (residual) capacity of the edges in P.
Cycle redistribution

return f

[Generalized](#page-26-0) assignment problems

```
Circulations
Lower bounds
Joint rounding
```
[Min cost Max](#page-98-0) Flow

Let $\mathcal{N} = (V, E, c, s, \text{S}, t)$ and let f be a flow in \mathcal{N} ,

Redistribute(C, f) b=bottleneck (C) for each $(u, v) \in C$ do if (u, v) is a forward edge then Increase $f(u, v)$ by b else

Decrease $f(v, u)$ by b

メロメ メ団 メメミメメミメ

 $2Q$

Cycle redistribution

[Generalized](#page-26-0) assignment problems

```
Circulations
Lower bounds
Joint rounding
```
[Min cost Max](#page-98-0) Flow

Let $\mathcal{N} = (V, E, c, s, \text{S}, t)$ and let f be a flow in \mathcal{N} , Redistribute(C, f) b=bottleneck (C) for each $(u, v) \in C$ do if (u, v) is a forward edge then Increase $f(u, v)$ by b else Decrease $f(v, u)$ by b s 2/2 Ω 2/3

return f

a b t 0/1 2/2 \mathcal{N} , f

つへへ

←ロ ▶ → 伊 ▶

Cycle redistribution

[Generalized](#page-26-0) assignment problems

[Circulations](#page-40-0) [Lower bounds](#page-65-0) [Joint rounding](#page-88-0)

[Min cost Max](#page-98-0) Flow

Let $\mathcal{N} = (V, E, c, s, \text{S}, t)$ and let f be a flow in \mathcal{N} , Redistribute(C, f) b=bottleneck (C) for each $(u, v) \in C$ do if (u, v) is a forward edge then Increase $f(u, v)$ by b else Decrease $f(v, u)$ by b return f s a b 2/2 Ω 0/1 2/2 2/3 N_f

t

[Edmonds](#page-1-0) Karp alg

[Generalized](#page-26-0) assignment problems

[Circulations](#page-40-0) [Lower bounds](#page-65-0) [Joint rounding](#page-88-0)

[Min cost Max](#page-98-0) Flow

Lemma

Let $f' = \text{Redistribute}(C, f)$, then f' is a flow in $\mathcal N$ and $|f'| = |f|.$

Proof.

We have to prove the two flow properties.

■ Capacity law

- Forward edges $(u, v) \in P$, we increase $f(u, v)$ by b, as $b \leq c(u, v) - f(u, v)$ then $f'(u, v) = f(u, v) + b \leq c(u, v)$.
- Backward edges $(u, v) \in P$ we decrease $f(v, u)$ by b, as $b \leq f(v, u), f'(v, u) = f(u, v) - b \geq 0.$

[Edmonds](#page-1-0) Karp alg

[Generalized](#page-26-0) assignment problems

[Circulations](#page-40-0) [Lower bounds](#page-65-0) [Joint rounding](#page-88-0)

[Min cost Max](#page-98-0) Flow

- Conservation law, $\forall v \in P \setminus \{s, t\}$ let u be the predecessor of v in P and let w be its successor.
- As the cycle is simple only the alterations due to (u, v) and (v, w) can change the flow that goes trough v. As we did for the algorithm Augment, for augmenting path, a case by case analysis shows that the conservation law is preserved.

へのへ

[Edmonds](#page-1-0) Karp alg

[Generalized](#page-26-0) assignment

[Circulations](#page-40-0) [Lower bounds](#page-65-0) [Joint rounding](#page-88-0)

[Min cost Max](#page-98-0) Flow

Now we have to prove that the value of the flow does not change. We have two cases:

メロメ メ御 メメ ミメメ ミメー

重

 $2Q$

Now we have to prove that the value of the flow does not change. We have two cases:

■ $s \notin C$. As Redistribute only changes the in/out flow of the vertices in C , the flow out of s is not changed. Therefore, $|f'| = |f|.$

∢ロ ▶ ∢ 御 ▶ ∢ 后 ▶ ∢ 后 ▶

つくい

[Edmonds](#page-1-0) Karp alg

[Generalized](#page-26-0) problems

[Circulations](#page-40-0) [Lower bounds](#page-65-0) [Joint rounding](#page-88-0)

[Min cost Max](#page-98-0) Flow

Now we have to prove that the value of the flow does not change. We have two cases:

- $s \notin C$. As Redistribute only changes the in/out flow of the vertices in C , the flow out of s is not changed. Therefore, $|f'| = |f|.$
- $s \in C$. As s has not incoming edges, any cycle in G_f containing s must involve a backward edge entering s and a forward edge out of s. Therefore, we reduce by b the flow in one edge out of s and increase by b another such edges. Again, $|f'| = |f|$.

[Edmonds](#page-1-0) Karp alg

[Generalized](#page-26-0) assignment problems

[Circulations](#page-40-0) [Lower bounds](#page-65-0) [Joint rounding](#page-88-0)

[Min cost Max](#page-98-0) Flow

メロメ メ御 メメ ミメメ ヨメ へのへ

EndProof

Redistribute: cost

[Edmonds](#page-1-0) Karp alg

[Generalized](#page-26-0)

[Circulations](#page-40-0) [Lower bounds](#page-65-0) [Joint rounding](#page-88-0)

[Min cost Max](#page-98-0) Flow

Lemma

Let $f' = \text{Redistribute}(C, f)$, then $\$(f') = \$(f) + b\$(f)$.

≮ロ ▶ ⊀ 御 ▶ ⊀ 君 ▶ ⊀ 君 ▶

重

 $2Q$

Redistribute: cost

[Generalized](#page-26-0) assignment problems

[Circulations](#page-40-0) [Lower bounds](#page-65-0) [Joint rounding](#page-88-0)

[Min cost Max](#page-98-0) Flow

Lemma

Let
$$
f' = \text{Redistribute}(C, f)
$$
, then $\$(f') = \$(f) + b\$(f) =$.

Proof.

The changes done by Redistribute are (1) subtract b units of flow from the backward edges in C and (2) add b units of flow to the forward edges in C. According to the definition of \mathcal{S}_f the total change in cost is given by $b\mathfrak{F}_f(C)$.

Min cost Max Flow

[Edmonds](#page-1-0) Karp alg

[Generalized](#page-26-0) assignment problems

[Circulations](#page-40-0) [Lower bounds](#page-65-0) [Joint rounding](#page-88-0)

[Min cost Max](#page-98-0) Flow

Theorem

f is a minimum cost maximum flow for $\mathcal{N} = (V, E, c, s, \text{ } \text{\^{S}, t)$ iff f is a maximum flow in $\mathcal{N} = (V, E, c, s, t)$ and the residual graph G_f has no negative cost cycles.

Proof.

If there is a negative cost cycle C , f has maximum value but not minimum cost as Redistribute(f, C) will provide a flow with maximum value and smaller cost.

Min cost Max Flow

[Edmonds](#page-1-0) Karp alg

[Generalized](#page-26-0) assignment problems

[Circulations](#page-40-0) [Lower bounds](#page-65-0) [Joint rounding](#page-88-0)

[Min cost Max](#page-98-0) Flow

If there is no negative cycle,

- We can compute the shortest distance $\delta(v)$ from s to every node v in G_f according to edge weight $\$_f.$
- As we have seen when discussing Johnson's algorithm, under the reduced cost $c(v,w) = \frac{6}{3}f(v,w) + \delta(v) - \delta(w)$, all edges in G_f have non-negative costs. This means that any change in f , cannot decrease the reduced cost of f .

4 n + 4 n + 4 =

By the path/cycle invariant of the reduced cost, any change in f cannot decrease its cost.

EndProof.

∽≏ເ

Cycle-canceling algorithm

- **[Edmonds](#page-1-0)** Karp alg
- [Generalized](#page-26-0)
- **[Circulations](#page-40-0)** [Lower bounds](#page-65-0) [Joint rounding](#page-88-0)

[Min cost Max](#page-98-0) Flow

メロメメ 倒 メメ きょくきょう 目 299

Cycle-canceling algorithm

[Edmonds](#page-1-0) Karp alg

[Generalized](#page-26-0) assignment

[Circulations](#page-40-0) [Lower bounds](#page-65-0) [Joint rounding](#page-88-0)

[Min cost Max](#page-98-0) Flow

Note that when $f' =$ Redistribute (F, C) for some negative cost cycle C in G_f , C does not form in $G_{f'}$.

メロメ メ御 メメ ミメメ ミメ

 $2Q$

€

Cycle-canceling algorithm

[Edmonds](#page-1-0) Karp alg

[Generalized](#page-26-0) assignment problems

[Circulations](#page-40-0) [Lower bounds](#page-65-0) [Joint rounding](#page-88-0)

[Min cost Max](#page-98-0) Flow

Note that when $f' =$ Redistribute (F, C) for some negative cost cycle C in G_f , C does not form in $G_{f'}$.

Morton Klein, A Primal Method for Minimal Cost Flows with Applications to the Assignment and Transportation Problems, Management Science, INFORMS, vol. 14(3), pages 205-220, November 1967.

Cycle Canceling $(G, s, t, c, \$)$ $f = \text{MaxFlow}(G, s, t)$ Compute G_f while there is a negative cost cycle C in G_f do $f = \text{Redistribute}(f, \mathcal{C}, \mathcal{G}_f)$ Compute G_f return f

Networks with integer capacities

Using the same arguments as for the Ford Fulkerson algorithm.

Lemma (Integrality invariant)

[Generalized](#page-26-0) assignment problems

[Edmonds](#page-1-0) Karp alg

[Circulations](#page-40-0) [Lower bounds](#page-65-0) [Joint rounding](#page-88-0)

[Min cost Max](#page-98-0) Flow

Let $\mathcal{N} = (V, E, c, \text{m}, s, t)$ where $c : E \to \mathbb{Z}^+$. At every iteration of the Cycle Canceling algorithm, the flow values $f(e)$ are integers.

Theorem (Integrality theorem)

Let $\mathcal{N} = (V, E, c, \$, s, s, t)$ where $c : E \rightarrow \mathbb{Z}^+$. There exists a min cost max-flow f^* such that $f^*(e)$ is an integer, for any $e \in E$.

Networks with integer capacities and costs

Lemma

[Generalized](#page-26-0) assignment problems

[Edmonds](#page-1-0) Karp alg

[Circulations](#page-40-0) [Lower bounds](#page-65-0) [Joint rounding](#page-88-0)

[Min cost Max](#page-98-0) Flow

Let $\mathcal{N} = (V, E, c, \texttt{\$}, s, t)$ where $c, \texttt{\$} : E \to \mathbb{Z}^{+}$. Let C be the min cut capacity, the Cycle Canceling algorithm terminates after finding at most C augmenting paths and after performing at most $$(C)$ redistribution calls.

Proof.

The value of the flow increases by ≥ 1 after each augmentation and the cost of a maximum flow after a redistribute call decreases at least by 1.

[Edmonds](#page-1-0) Karp alg

[Generalized](#page-26-0) assignment

[Circulations](#page-40-0) [Lower bounds](#page-65-0) [Joint rounding](#page-88-0)

[Min cost Max](#page-98-0) Flow

Computing the MaxFlow f^* takes $O(|f^*|(n+m))$.

メロト メタト メミト メミト

重

 $2Q$

[Edmonds](#page-1-0) Karp alg

[Generalized](#page-26-0) assignment problems

[Circulations](#page-40-0) [Lower bounds](#page-65-0) [Joint rounding](#page-88-0)

[Min cost Max](#page-98-0) Flow

Computing the MaxFlow f^* takes $O(|f^*|(n+m))$. For the second part of the Cycle Canceling algorithm: $\mathcal{L}_{\mathcal{A}}$

 $2Q$

メロメ メ団 メメミメメミメ

[Edmonds](#page-1-0) Karp alg

[Generalized](#page-26-0) assignment problems

[Circulations](#page-40-0) [Lower bounds](#page-65-0) [Joint rounding](#page-88-0)

[Min cost Max](#page-98-0) Flow

Computing the MaxFlow f^* takes $O(|f^*|(n+m))$.

For the second part of the Cycle Canceling algorithm: $\mathcal{L}_{\mathcal{A}}$

- Constructing G_f , takes $O(m)$ time.
- \blacksquare O(nm) time to decide if G_f has a negative cycle and if so computing one (use Bellman-Ford algorithm).

∢ロ ▶ ∢ 御 ▶ ∢ 后 ▶ ∢ 后 ▶

へのへ

A call to Redistribute requires $O(m)$ steps

[Edmonds](#page-1-0) Karp alg

[Generalized](#page-26-0) assignment problems

[Circulations](#page-40-0) [Lower bounds](#page-65-0) [Joint rounding](#page-88-0)

[Min cost Max](#page-98-0) Flow

Г

Computing the MaxFlow f^* takes $O(|f^*|(n+m))$.

For the second part of the Cycle Canceling algorithm:

- Constructing G_f , takes $O(m)$ time.
- \blacksquare O(nm) time to decide if G_f has a negative cycle and if so computing one (use Bellman-Ford algorithm).

∢ロ ▶ ∢ 御 ▶ ∢ 后 ▶ ∢ 后 ▶

へのへ

A call to Redistribute requires $O(m)$ steps

Let
$$
C = \max_{e \in E} c(e)
$$
 and $K = \max_{e \in E} \S(e)$,
 $\S(f^*) \leq CKm$

[Edmonds](#page-1-0) Karp alg

[Generalized](#page-26-0) assignment problems

[Circulations](#page-40-0) [Lower bounds](#page-65-0) [Joint rounding](#page-88-0)

[Min cost Max](#page-98-0) Flow

Computing the MaxFlow f^* takes $O(|f^*|(n+m))$.

For the second part of the Cycle Canceling algorithm: $\mathcal{L}_{\mathcal{A}}$

- Constructing G_f , takes $O(m)$ time.
- \blacksquare O(nm) time to decide if G_f has a negative cycle and if so computing one (use Bellman-Ford algorithm).

∢ロ ▶ ∢ 御 ▶ ∢ 后 ▶ ∢ 后 ▶

へのへ

A call to Redistribute requires $O(m)$ steps

Let
$$
C = \max_{e \in E} c(e)
$$
 and $K = \max_{e \in E} \S(e)$,
 $\S(f^*) \leq CKm$

Total running time is $O(|f^*|(n+m) + CKnm)$

[Edmonds](#page-1-0) Karp alg

[Generalized](#page-26-0) assignment problems

[Circulations](#page-40-0) [Lower bounds](#page-65-0) [Joint rounding](#page-88-0)

[Min cost Max](#page-98-0) Flow

- Computing the MaxFlow f^* takes $O(|f^*|(n+m))$.
- For the second part of the Cycle Canceling algorithm: $\mathcal{L}^{\mathcal{A}}$
	- Constructing G_f , takes $O(m)$ time.
	- \blacksquare O(nm) time to decide if G_f has a negative cycle and if so computing one (use Bellman-Ford algorithm).
	- A call to Redistribute requires $O(m)$ steps

Let
$$
C = \max_{e \in E} c(e)
$$
 and $K = \max_{e \in E} \S(e)$,
 $\S(f^*) \leq CKm$

- Total running time is $O(|f^*|(n+m) + CKnm)$
- **Thus, we have a pseudo polynomial algorithm.**

[Edmonds](#page-1-0) Karp alg

[Generalized](#page-26-0) assignment problems

[Circulations](#page-40-0) [Lower bounds](#page-65-0) [Joint rounding](#page-88-0)

[Min cost Max](#page-98-0) Flow

- Computing the MaxFlow f^* takes $O(|f^*|(n+m))$.
- For the second part of the Cycle Canceling algorithm: $\mathcal{L}^{\mathcal{A}}$
	- Constructing G_f , takes $O(m)$ time.
	- \blacksquare O(nm) time to decide if G_f has a negative cycle and if so computing one (use Bellman-Ford algorithm).
	- A call to Redistribute requires $O(m)$ steps

Let
$$
C = \max_{e \in E} c(e)
$$
 and $K = \max_{e \in E} \S(e)$,
 $\S(f^*) \leq CKm$

- Total running time is $O(|f^*|(n+m) + CKnm)$
- **Thus, we have a pseudo polynomial algorithm.**

[Edmonds](#page-1-0) Karp alg

[Generalized](#page-26-0) assignment

[Circulations](#page-40-0) [Lower bounds](#page-65-0) [Joint rounding](#page-88-0)

[Min cost Max](#page-98-0) Flow

Like Ford-Fulkerson algorithm, more careful choices of which cycle to cancel lead to more efficient algorithms.

メロト メタト メミト メミト

重

 $2Q$

- **[Edmonds](#page-1-0)** Karp alg
- [Generalized](#page-26-0) assignment problems
- **[Circulations](#page-40-0)** [Lower bounds](#page-65-0) [Joint rounding](#page-88-0)

[Min cost Max](#page-98-0) Flow

- Like Ford-Fulkerson algorithm, more careful choices of which cycle to cancel lead to more efficient algorithms.
- In 1980, Goldberg and Tarjan developed an algorithm that cancels the minimum-mean cycle, the cycle whose average cost per edge is smallest. A clever implementation of the algorithm achieves running time $O(nm^2 \log V)$

∢ロ ▶ ∢ 御 ▶ ∢ 后 ▶ ∢ 后 ▶

へのへ

[Edmonds](#page-1-0) Karp alg

[Generalized](#page-26-0) assignment problems

[Circulations](#page-40-0) [Lower bounds](#page-65-0) [Joint rounding](#page-88-0)

[Min cost Max](#page-98-0) Flow

- Like Ford-Fulkerson algorithm, more careful choices of which cycle to cancel lead to more efficient algorithms.
- In 1980, Goldberg and Tarjan developed an algorithm that cancels the minimum-mean cycle, the cycle whose average cost per edge is smallest. A clever implementation of the algorithm achieves running time $O(nm^2 \log V)$
- Combining Edmonds-Karp algorithm with Goldberg and Tarjan's, we get a polynomial time algorithm solving the Min cost Maximum flow problem.

[Edmonds](#page-1-0) Karp alg

[Generalized](#page-26-0) assignment problems

[Circulations](#page-40-0) [Lower bounds](#page-65-0) [Joint rounding](#page-88-0)

[Min cost Max](#page-98-0) Flow

- Like Ford-Fulkerson algorithm, more careful choices of which cycle to cancel lead to more efficient algorithms.
- In 1980, Goldberg and Tarjan developed an algorithm that cancels the minimum-mean cycle, the cycle whose average cost per edge is smallest. A clever implementation of the algorithm achieves running time $O(nm^2 \log V)$
- Combining Edmonds-Karp algorithm with Goldberg and Tarjan's, we get a polynomial time algorithm solving the Min cost Maximum flow problem.
- The, Chen, Kyng, Liu, Peng, Gutenberg, Sachdeva (2022) $O(m^{1+o(1)})$ solves also the Min cost Maximum flow algorithm.

Min cost circulations

- **[Edmonds](#page-1-0)** Karp alg
- [Generalized](#page-26-0) assignment problems
- **[Circulations](#page-40-0)** [Lower bounds](#page-65-0) [Joint rounding](#page-88-0)

[Min cost Max](#page-98-0) Flow

- The Cycle Canceling algorithm can be extended to compute min cost circulations in flow networks with demands and lower bounds, provided a circulation exists.
- \blacksquare The algorithms, have the same asymptotic cost as the ones for the minimum cost maximum flow problem

4 n + 4 n + 4 =

へのへ