

Max-flow and min-cut problems

Max Flow and
Min Cut

Properties of
flows and cuts

Residual
graph

Augmenting
path

MaxFlow
MinCut Thm

Ford
Fulkerson alg



- 1 Max Flow and Min Cut
- 2 Properties of flows and cuts
- 3 Residual graph
- 4 Augmenting path
- 5 MaxFlow MinCut Thm
- 6 Ford Fulkerson alg

Flow Network

Max Flow and
Min Cut

Properties of
flows and cuts

Residual
graph


Augmenting
path

MaxFlow
MinCut Thm

Ford
Fulkerson alg

A network $\mathcal{N} = (V, E, c, s, t)$ is formed by

- a digraph $G = (V, E)$,
- a source vertex $s \in V$
- a sink vertex $t \in V$,
- and edge capacities $c : E \rightarrow \mathbb{R}^+$



A flow in a network

Max Flow and
Min Cut

Properties of
flows and cuts

Residual
graph

Augmenting
path

MaxFlow
MinCut Thm

Ford
Fulkerson alg


Given a network $\mathcal{N} = (V, E, c, s, t)$

A **Flow** is an assignment $f : E \rightarrow \mathbb{R}^+ \cup \{0\}$ that follows the **Kirchoff's laws**:

- $\forall (u, v) \in E, 0 \leq f(u, v) \leq c(u, v),$
- (Flow conservation) $\forall v \in V - \{s, t\},$
 $\sum_{u \in V} f(u, v) = \sum_{z \in V} f(v, z)$

The **value of a flow** f is

$$|f| = \sum_{v \in V} f(s, v) = f(s, V) = f(V, t).$$



$f(e)/c(e)$

A flow in a network

Max Flow and
Min Cut

Properties of
flows and cuts

Residual
graph

Augmenting
path

MaxFlow
MinCut Thm

Ford
Fulkerson alg


Given a network $\mathcal{N} = (V, E, c, s, t)$

A **Flow** is an assignment $f : E \rightarrow \mathbb{R}^+ \cup \{0\}$ that follows the **Kirchoff's laws**:

- $\forall (u, v) \in E, 0 \leq f(u, v) \leq c(u, v),$
- (Flow conservation) $\forall v \in V - \{s, t\},$
 $\sum_{u \in V} f(u, v) = \sum_{z \in V} f(v, z)$

The **value of a flow** f is

$$|f| = \sum_{v \in V} f(s, v) = f(s, V) = f(V, t).$$



$$f(e)/c(e)$$

with value 3.

A flow in a network

Max Flow and
Min Cut

Properties of
flows and cuts

Residual
graph

Augmenting
path

MaxFlow
MinCut Thm

Ford
Fulkerson alg


Given a network $\mathcal{N} = (V, E, c, s, t)$

A **Flow** is an assignment $f : E \rightarrow \mathbb{R}^+ \cup \{0\}$ that follows the **Kirchoff's laws**:

- $\forall (u, v) \in E, 0 \leq f(u, v) \leq c(u, v),$
- (Flow conservation) $\forall v \in V - \{s, t\},$
$$\sum_{u \in V} f(u, v) = \sum_{z \in V} f(v, z)$$

The **value of a flow** f is

$$|f| = \sum_{v \in V} f(s, v) = f(s, V) = f(V, t).$$



saturated

The Maximum flow problem

INPUT: A network $\mathcal{N} = (V, E, c, s, t,)$

QUESTION: Find a flow of maximum value on \mathcal{N} .

Max Flow and
Min Cut

Properties of
flows and cuts

Residual
graph

Augmenting
path


MaxFlow
MinCut Thm

Ford
Fulkerson alg

The Maximum flow problem

INPUT: A network $\mathcal{N} = (V, E, c, s, t,)$

QUESTION: Find a flow of maximum value on \mathcal{N} .



Max Flow and
Min Cut

Properties of
flows and cuts

Residual
graph

Augmenting
path


MaxFlow
MinCut Thm

Ford
Fulkerson alg

The Maximum flow problem

INPUT: A network $\mathcal{N} = (V, E, c, s, t,)$

QUESTION: Find a flow of maximum value on \mathcal{N} .



The value of the flow is **7** = 4 + 1 + 2 = 5 + 2.

As t cannot receive more flow, this flow is a **maximum flow**.

Max Flow and
Min Cut

Properties of
flows and cuts

Residual
graph

Augmenting
path

MaxFlow
MinCut Thm


Ford
Fulkerson alg

The (s, t) -cuts

Given $\mathcal{N} = (V, E, c, s, t)$ a **(s, t) -cut** is a partition of $V = S \cup T$ ($S \cap T = \emptyset$), with $s \in S$ and $t \in T$.

The **capacity** of a cut (S, T) is the sum of weights **leaving** S , i.e.,

$$c(S, T) = \sum_{u \in S} \sum_{v \in T} c(u, v)$$



$$S = \{s, c, d\}$$

$$T = \{a, b, e, t\}$$

$$c(S, T) = 19$$

$$(4 + 5) + 5 + (3 + 2)$$

Max Flow and
Min Cut

Properties of
flows and cuts

Residual
graph

Augmenting
path

MaxFlow
MinCut Thm


Ford
Fulkerson alg

The (s, t) -cuts

Given $\mathcal{N} = (V, E, c, s, t)$ a **(s, t) -cut** is a partition of $V = S \cup T$ ($S \cap T = \emptyset$), with $s \in S$ and $t \in T$.

The flow across the cut:

$$f(S, T) = \sum_{u \in S} \sum_{v \in T} f(u, v) - \sum_{v \in T} \sum_{u \in S} f(v, u).$$



$$\begin{aligned} S &= \{s, c, d\} \\ T &= \{a, b, e, t\} \\ c(S, T) &= 19 \\ f(S, T) &= 10 - 3 = 7 \end{aligned}$$

Max Flow and
Min Cut

Properties of
flows and cuts

Residual
graph

Augmenting
path

MaxFlow
MinCut Thm


Ford
Fulkerson alg

The (s, t) -cuts

Given $\mathcal{N} = (V, E, c, s, t)$ a **(s, t) -cut** is a partition of $V = S \cup T$ ($S \cap T = \emptyset$), with $s \in S$ and $t \in T$.

The flow across the cut:

$$f(S, T) = \sum_{u \in S} \sum_{v \in T} f(u, v) - \sum_{v \in T} \sum_{u \in S} f(v, u).$$



$$\begin{aligned} S &= \{s, c, d\} \\ T &= \{a, b, e, t\} \\ c(S, T) &= 19 \\ f(S, T) &= 10 - 3 = 7 \end{aligned}$$

Due to the capacity constrain: $f(S, T) \leq c(S, T)$

Max Flow and
Min Cut

Properties of
flows and cuts

Residual
graph

Augmenting
path

MaxFlow
MinCut Thm


Ford
Fulkerson alg

Another (s, t) -cut

Given $\mathcal{N} = (V, E, c, s, t)$ a **(s, t) -cut** is a partition of $V = S \cup T$ ($S \cap T = \emptyset$), with $s \in S$ and $t \in T$.

The flow across the cut:

$$f(S, T) = \sum_{u \in S} \sum_{v \in T} f(u, v) - \sum_{v \in T} \sum_{u \in S} f(v, u).$$



$$S = \{s, a, c\}$$

$$T = \{b, d, e, t\}$$

$$c(S, T) = 12$$

$$f(S, T) = 8 - 1 = 7$$

Max Flow and
Min Cut

Properties of
flows and cuts

Residual
graph

Augmenting
path

MaxFlow
MinCut Thm

Ford
Fulkerson alg

The Minimum Cut problem

Max Flow and
Min Cut

Properties of
flows and cuts

Residual
graph


Augmenting
path

MaxFlow
MinCut Thm

Ford
Fulkerson alg

INPUT: A network $\mathcal{N} = (V, E, c, s, t,)$

QUESTION: Find a (s, t) -cut of minimum capacity in \mathcal{N} .



MinCut

$S = \{s, a, b, c, d, e\}$

$T = \{t\}$

$c(S, T) = 7$

Changing weights effect on min cuts

Given a network $\mathcal{N} = (V, E, s, t, c)$ assume that (S, T) is a min (s, t) -cut.

Max Flow and
Min Cut

Properties of
flows and cuts

Residual
graph

Augmenting
path

MaxFlow
MinCut Thm

Ford
Fulkerson alg

Changing weights effect on min cuts

Max Flow and
Min Cut

Properties of
flows and cuts

Residual
graph



Augmenting
path

MaxFlow
MinCut Thm

Ford
Fulkerson alg

Given a network $\mathcal{N} = (V, E, s, t, c)$ assume that (S, T) is a min (s, t) -cut.

If we change the input by adding $c > 0$ to the capacity of **every edge**, then it may happen that (S, T) is not longer a min (s, t) -cut.



Changing weights effect on Min-Cut and Max-Flow

Max Flow and
Min Cut

Properties of
flows and cuts

Residual
graph

Augmenting
path

MaxFlow
MinCut Thm

Ford
Fulkerson alg

Given a network $\mathcal{N} = (V, E, s, t, c)$.

Changing weights effect on Min-Cut and Max-Flow

Max Flow and
Min Cut

Properties of
flows and cuts

Residual
graph



Augmenting
path

MaxFlow
MinCut Thm

Ford
Fulkerson alg

Given a network $\mathcal{N} = (V, E, s, t, c)$.

If we change the network by multiplying by $c > 1$ the capacity of every edge, the capacity of any (s, t) -cut in the new network is c times its capacity in the original network.



- 1 Max Flow and Min Cut
- 2 Properties of flows and cuts**
- 3 Residual graph
- 4 Augmenting path
- 5 MaxFlow MinCut Thm
- 6 Ford Fulkerson alg

Notation

Let $\mathcal{N} = (V, E, s, t, c)$ and f a flow in \mathcal{N}

For $v \in V$, $U \subseteq V$ and $v \notin U$.

- $f(v, U)$ flow $v \rightarrow U$ i.e. $f(v, U) = \sum_{u \in U} f(v, u)$,
- $f(U, v)$ flow $U \rightarrow v$ i.e. $f(U, v) = \sum_{u \in U} f(u, v)$,

For a (s, t) -cut (S, T) and $v \in S$

- $S' = S \setminus \{v\}$ and $T' = T \cup \{v\}$
- $f_{-v}(S, T) = \sum_{u \in S'} \sum_{w \in T} f(u, w) - \sum_{w \in T} \sum_{u \in S'} f(w, u)$
i.e, the contribution to $f(S, T)$ from edges not incident with v .

Max Flow and
Min Cut

Properties of
flows and cuts

Residual
graph

Augmenting
path

MaxFlow
MinCut Thm

Ford
Fulkerson alg

Flow conservation on (s, t) -cuts

Max Flow and
Min Cut

Properties of
flows and cuts

Residual
graph

Augmenting
path

MaxFlow
MinCut Thm

Ford
Fulkerson alg

Theorem

Let $\mathcal{N} = (V, E, s, t, c)$ and f a flow in \mathcal{N} . For any (s, t) -cut (S, T) , $f(S, T) = |f|$.

Proof (Induction on $|S|$)

- If $S = \{s\}$ then, by definition, $f(S, T) = |f|$.

Flow conservation on (s, t) -cuts

Max Flow and
Min Cut

Properties of
flows and cuts

Residual
graph

Augmenting
path

MaxFlow
MinCut Thm

Ford
Fulkerson alg

Theorem


Let $\mathcal{N} = (V, E, s, t, c)$ and f a flow in \mathcal{N} . For any (s, t) -cut (S, T) , $f(S, T) = |f|$.

Proof (Induction on $|S|$)

- If $S = \{s\}$ then, by definition, $f(S, T) = |f|$.
- Assume it is true for $S' = S - \{v\}$ and $T' = T \cup \{v\}$, i.e. $f(S', T') = |f|$.

Flow conservation on (s, t) -cuts

Proof (cont.) (Induction on $|S|$)



- IH: $f(S', T') = |f|$.
- Then, $f(S, T) = f_{-v}(S, T) + f(v, T) - f(T, v)$.
- But, $f(S', T') = f_{-v}(S, T) + f(S', v) - f(v, S')$ as $v \in T'$
- By flow conservation,
$$f(S', v) + f(T, v) = f(v, S') + f(v, T)$$
- So, $f(S', v) - f(v, S') = f(v, T) - f(T, v)$
- Therefore, $f(S', T') = f(S, T) = |f|$



Max Flow and
Min Cut

Properties of
flows and cuts

**Residual
graph**

Augmenting
path

MaxFlow
MinCut Thm



Ford
Fulkerson alg

- 1 Max Flow and Min Cut
- 2 Properties of flows and cuts
- 3 Residual graph**
- 4 Augmenting path
- 5 MaxFlow MinCut Thm
- 6 Ford Fulkerson alg

Residual graph

Given a network $\mathcal{N} = (V, E, s, t, c)$ together with a **flow** f .
The **residual graph**, $(G_f = (V, E_f, c_f))$ is a weighted digraph on the same vertex set and with edge set:



- if $c(u, v) - f(u, v) > 0$, then $(u, v) \in E_f$ and $c_f(u, v) = c(u, v) - f(u, v) > 0$ (**forward edges**)



Residual graph

Given a network $\mathcal{N} = (V, E, s, t, c)$ together with a **flow** f on it, the **residual graph**, $(G_f = (V, E_f, c_f))$ is a weighted digraph on the same vertex set and with edge set:

- if $f(u, v) > 0$, then $(v, u) \in E_f$ and $c_f(v, u) = f(u, v)$ (**backward edges**).



Max Flow and
Min Cut

Properties of
flows and cuts

Residual
graph

Augmenting
path



MaxFlow
MinCut Thm

Ford
Fulkerson alg

Residual graph

Given a network $\mathcal{N} = (V, E, s, t, c)$ together with a **flow f** on it, the **residual graph**, $(G_f = (V, E_f, c_f))$ is a weighted digraph on the same vertex set and with edge set:

- if $c(u, v) - f(u, v) > 0$, then $(u, v) \in E_f$ and $c_f(u, v) = c(u, v) - f(u, v) > 0$ (**forward edges**)
- if $f(u, v) > 0$, then $(v, u) \in E_f$ and $c_f(v, u) = f(u, v)$ (**backward edges**).



Max Flow and
Min Cut

Properties of
flows and cuts



Residual
graph

Augmenting
path

MaxFlow
MinCut Thm



Ford
Fulkerson alg

Residual graph



- Notice that, if $c(u, v) = f(u, v)$, then there is only a backward edge.
- c_f are called the residual capacity.

Residual graph



- **forward edges:** There remains capacity to push more flow through this edge.
- **backward edges:** there are units of flow that can be redirected through other links.

Max Flow and
Min Cut

Properties of
flows and cuts

Residual
graph

**Augmenting
path**



MaxFlow
MinCut Thm

Ford
Fulkerson alg

- 1 Max Flow and Min Cut
- 2 Properties of flows and cuts
- 3 Residual graph
- 4 Augmenting path**
- 5 MaxFlow MinCut Thm
- 6 Ford Fulkerson alg

Augmenting paths



Let $\mathcal{N} = (V, E, c, s, t)$ and let f be a flow in \mathcal{N} ,



- An **augmenting path** P is any **simple** path P in G_f from s to t

Augmenting paths



Let $\mathcal{N} = (V, E, c, s, t)$ and let f be a flow in \mathcal{N} ,



- An **augmenting path** P is any **simple path** P in G_f from s to t . P might have forward and backward edges.

Augmenting paths


Let $\mathcal{N} = (V, E, c, s, t)$ and let f be a flow in \mathcal{N} ,



- An **augmenting path** P is any **simple** path P in G_f from s to t . P might have forward and backward edges.
- For an augmenting path P in G_f , the **bottleneck**, $b(P)$, is the minimum (residual) capacity of the edges in P . In the example, for $P = (s, b, a, t)$, $b(P) = 1$.

Augmenting paths: increasing the flow

```
Augment( $P, f$ )  
   $b = \text{bottleneck}(P)$   
  for each  $(u, v) \in P$  do  
    if  $(u, v)$  is a forward edge then  
      Increase  $f(u, v)$  by  $b$   
    else  
      Decrease  $f(v, u)$  by  $b$   
  return  $f$ 
```



Max Flow and
Min Cut

Properties of
flows and cuts

Residual
graph



Augmenting
path

MaxFlow
MinCut Thm

Ford
Fulkerson alg

Augmenting paths: increasing the flow

Augment(P, f)
b=bottleneck (P)
for each $(u, v) \in P$ **do**
 if (u, v) is a forward edge **then**
 Increase $f(u, v)$ by b
 else
 Decrease $f(v, u)$ by b
return f



$G_f, P = (s, b, a, t), b(P) = 1$

Max Flow and
Min Cut

Properties of
flows and cuts

Residual
graph




Augmenting
path

MaxFlow
MinCut Thm

Ford
Fulkerson alg

Augmenting paths: increasing the flow

Augment(P, f)
b=bottleneck (P)
for each $(u, v) \in P$ **do**
 if (u, v) is a forward edge **then**
 Increase $f(u, v)$ by b
 else
 Decrease $f(v, u)$ by b
return f



Augmenting paths: increasing the flow

Max Flow and
Min Cut

Properties of
flows and cuts

Residual
graph

Augmenting
path

MaxFlow
MinCut Thm

Ford
Fulkerson alg

Lemma

Let $f' = \text{Augment}(P, f)$, then f' is a flow in G .

Proof: We have to prove the two flow properties.

■ Capacity law

- Forward edges $(u, v) \in P$, we increase $f(u, v)$ by b , as $b \leq c(u, v) - f(u, v)$ then $f'(u, v) = f(u, v) + b \leq c(u, v)$.
- Backward edges $(u, v) \in P$ we decrease $f(v, u)$ by b , as $b \leq f(v, u)$, $f'(v, u) = f(v, u) - b \geq 0$.

Augmenting paths: increasing the flow

Lemma

Let $f' = \text{Augment}(P, f)$, then f' is a flow in G .

Proof: We have to prove the two flow properties.

- **Conservation law**, $\forall v \in P \setminus \{s, t\}$ let u be the predecessor of v in P and let w be its successor.
- As the path is simple only the alterations due to (u, v) and (v, w) can change the flow that goes through v . We have four cases:
 - (u, v) and (v, w) are backward edges, the flow in (v, u) and (w, v) is decremented by b . As one is incoming and the other outgoing the total balance is 0.
 - (u, v) and (v, w) are forward edges, the flow in (u, v) and (v, w) is incremented by b . As one is incoming and the other outgoing the total balance is 0.

Augmenting paths: increasing the flow

Lemma

Let $f' = \text{Augment}(P, f)$, then f' is a flow in G .

Proof: We have to prove the two flow properties.

- **Conservation law**, $\forall v \in P \setminus \{s, t\}$ let u be the predecessor of v in P and let w be its successor.
- As the path is simple only the alterations due to (u, v) and (v, w) can change the flow that goes through v . We have three cases:
 - (u, v) is forward and (v, w) is backward, the flow in (u, v) is incremented by b and the flow in (w, v) is decremented by b . As both are incoming, the total balance is 0.
 - (u, v) is backward and (v, w) is forward, the flow in (v, w) is incremented by b and the flow in (v, u) is decremented by b . As both are outgoing, the total balance is 0.

Augmenting paths: incrementing the flow

Max Flow and
Min Cut

Properties of
flows and cuts

Residual
graph

**Augmenting
path**

MaxFlow
MinCut Thm

Ford
Fulkerson alg

Lemma

Consider $f' = \text{Augment}(P, f)$, then $|f'| > |f|$.

Proof: Let P be the augmenting path in G_f . The first edge $e \in P$ leaves s , and as G has no incoming edges to s , e is a forward edge. Moreover P is simple \Rightarrow never returns to s . Therefore, the value of the flow increases in edge e by b units.

□

Max Flow and
Min Cut

Properties of
flows and cuts

Residual
graph

Augmenting
path

**MaxFlow
MinCut Thm**

Ford
Fulkerson alg

- 1 Max Flow and Min Cut
- 2 Properties of flows and cuts
- 3 Residual graph
- 4 Augmenting path
- 5 MaxFlow MinCut Thm**
- 6 Ford Fulkerson alg

Max-Flow Min-Cut theorem

Max Flow and
Min Cut

Properties of
flows and cuts

Residual
graph

Augmenting
path

MaxFlow
MinCut Thm

Ford
Fulkerson alg

Ford and Fulkerson (1954); Peter Elias, Amiel Feinstein and Claude Shannon (1956) (in framework of information-theory).

Theorem

For any $\mathcal{N}(G, s, t, c)$, the maximum of the flow value is equal to the minimum of the (S, T) -cut capacities.

$$\max_f \{|f|\} = \min_{(S, T)} \{c(S, T)\}.$$

Max-Flow Min-Cut theorem: Proof

Proof:

- Let f^* be a flow with maximum value, $|f^*| = \max_f \{|f|\}$

Max Flow and
Min Cut

Properties of
flows and cuts

Residual
graph

Augmenting
path

**MaxFlow
MinCut Thm**

Ford
Fulkerson alg

Max-Flow Min-Cut theorem: Proof

Proof:

- Let f^* be a flow with maximum value, $|f^*| = \max_f \{|f|\}$
- For any (s, t) -cut (S, T) , $f^*(S, T) \leq c(S, T)$.

Max Flow and
Min Cut

Properties of
flows and cuts

Residual
graph

Augmenting
path

**MaxFlow
MinCut Thm**

Ford
Fulkerson alg

Max-Flow Min-Cut theorem: Proof

Proof:

- Let f^* be a flow with maximum value, $|f^*| = \max_f \{|f|\}$
- For any (s, t) -cut (S, T) , $f^*(S, T) \leq c(S, T)$.
- G_{f^*} has no augmenting path. So, if $S_s = \{v \in V \mid \exists s \rightsquigarrow v \text{ in } G_{f^*}\}$, then $(S_s, V - \{S_s\})$ is a (s, t) -cut.

Max Flow and
Min Cut

Properties of
flows and cuts

Residual
graph

Augmenting
path

MaxFlow
MinCut Thm

Ford
Fulkerson alg

Max-Flow Min-Cut theorem: Proof

Proof:

- Let f^* be a flow with maximum value, $|f^*| = \max_f \{|f|\}$
- For any (s, t) -cut (S, T) , $f^*(S, T) \leq c(S, T)$.
- G_{f^*} has no augmenting path. So, if $S_s = \{v \in V \mid \exists s \rightsquigarrow v \text{ in } G_{f^*}\}$, then $(S_s, V - \{S_s\})$ is a (s, t) -cut.
- For $e = (u, v) \in E$ with $u \in S_s$ and $v \notin S_s$, $(u, v) \notin E(G_{f^*})$, therefore $f^*(u, v) = c(u, v)$,

Max Flow and
Min Cut

Properties of
flows and cuts

Residual
graph

Augmenting
path

MaxFlow
MinCut Thm

Ford
Fulkerson alg

Max-Flow Min-Cut theorem: Proof

Proof:

- Let f^* be a flow with maximum value, $|f^*| = \max_f \{|f|\}$
- For any (s, t) -cut (S, T) , $f^*(S, T) \leq c(S, T)$.
- G_{f^*} has no augmenting path. So, if $S_s = \{v \in V \mid \exists s \rightsquigarrow v \text{ in } G_{f^*}\}$, then $(S_s, V - \{S_s\})$ is a (s, t) -cut.
- For $e = (u, v) \in E$ with $u \in S_s$ and $v \notin S_s$, $(u, v) \notin E(G_{f^*})$, therefore $f^*(u, v) = c(u, v)$,
- Then, $c(S_s, V - \{S_s\}) = f^*(S_s, V - \{S_s\}) = |f^*|$

Max Flow and
Min Cut

Properties of
flows and cuts

Residual
graph

Augmenting
path

MaxFlow
MinCut Thm

Ford
Fulkerson alg

Max-Flow Min-Cut theorem: Proof

Proof:

- Let f^* be a flow with maximum value, $|f^*| = \max_f \{|f|\}$
- For any (s, t) -cut (S, T) , $f^*(S, T) \leq c(S, T)$.
- G_{f^*} has no augmenting path. So, if $S_s = \{v \in V \mid \exists s \rightsquigarrow v \text{ in } G_{f^*}\}$, then $(S_s, V - \{S_s\})$ is a (s, t) -cut.
- For $e = (u, v) \in E$ with $u \in S_s$ and $v \notin S_s$, $(u, v) \notin E(G_{f^*})$, therefore $f^*(u, v) = c(u, v)$,
- Then, $c(S_s, V - \{S_s\}) = f^*(S_s, V - \{S_s\}) = |f^*|$
- $(S_s, V - \{S_s\})$ is a minimum capacity (s, t) -cut in G .



Max Flow and
Min Cut

Properties of
flows and cuts

Residual
graph

Augmenting
path

MaxFlow
MinCut Thm

Ford
Fulkerson alg

- 1 Max Flow and Min Cut
- 2 Properties of flows and cuts
- 3 Residual graph
- 4 Augmenting path
- 5 MaxFlow MinCut Thm
- 6 Ford Fulkerson alg**

Ford-Fulkerson algorithm

Max Flow and
Min Cut

Properties of
flows and cuts


Residual
graph

Augmenting
path

MaxFlow
MinCut Thm


Ford
Fulkerson alg

L.R. Ford, D.R. Fulkerson:
*Maximal flow through a
network*. Canadian J. of Math.
1956.



```
Ford-Fulkerson( $G, s, t, c$ )  
for all  $(u, v) \in E$  set  $f(u, v) = 0$   
 $G_f = G$   
while there is an  $(s, t)$  path  $P$  in  $G_f$  do  
     $f = \text{Augment}(P, G_f)$   
    Compute  $G_f$   
return  $f$ 
```

FF algorithm example



Max Flow and
Min Cut

Properties of
flows and cuts



Residual
graph

Augmenting
path

MaxFlow
MinCut Thm

Ford
Fulkerson alg

FF algorithm example



Max Flow and
Min Cut

Properties of
flows and cuts

Residual
graph

Augmenting
path

MaxFlow
MinCut Thm

Ford
Fulkerson alg

FF algorithm example

Max Flow and
Min Cut


Properties of
flows and cuts

Residual
graph


Augmenting
path

MaxFlow
MinCut Thm


Ford
Fulkerson alg



Initial f_0



G_{f_0}



f_1

FF algorithm example

Max Flow and
Min Cut




Properties of
flows and cuts

Residual
graph

Augmenting
path

MaxFlow
MinCut Thm

Ford
Fulkerson alg



FF example

Max Flow and
Min Cut



Properties of
flows and cuts

Residual
graph

Augmenting
path

MaxFlow
MinCut Thm

Ford
Fulkerson alg



FF example

Max Flow and
Min Cut




Properties of
flows and cuts

Residual
graph

Augmenting
path

MaxFlow
MinCut Thm

Ford
Fulkerson alg



FF example

Max Flow and
Min Cut





Properties of
flows and cuts

Residual
graph

Augmenting
path

MaxFlow
MinCut Thm

Ford
Fulkerson alg



FF example

Max Flow and
Min Cut



Properties of
flows and cuts

Residual
graph

Augmenting
path

MaxFlow
MinCut Thm

Ford
Fulkerson alg



FF example

Max Flow and
Min Cut




Properties of
flows and cuts

Residual
graph

Augmenting
path

MaxFlow
MinCut Thm

Ford
Fulkerson alg



FF example

Max Flow and
Min Cut





Properties of
flows and cuts

Residual
graph

Augmenting
path

MaxFlow
MinCut Thm

Ford
Fulkerson alg



FF example

Max Flow and
Min Cut


Properties of
flows and cuts

Residual
graph


Augmenting
path

MaxFlow
MinCut Thm

Ford
Fulkerson alg



Flow with max value



$\{s\}, \{a, b, t\}$ is a min (s, t) -cut

Correctness of Ford-Fulkerson

Max Flow and
Min Cut

Properties of
flows and cuts

Residual
graph

Augmenting
path

MaxFlow
MinCut Thm

**Ford
Fulkerson alg**

Consequence of the Max-flow min-cut theorem.

Theorem

The flow returned by Ford-Fulkerson is the max-flow.

Networks with integer capacities

Lemma (*Integrality invariant*)

Let $\mathcal{N} = (V, E, c, s, t)$ where $c : E \rightarrow \mathbb{Z}^+$. At every iteration of the Ford-Fulkerson algorithm, the flow values $f(e)$ are integers.

Proof: (induction)

- The statement is true for the initial flow (all zeroes).
- Inductive Hypothesis: The statement is true after j iterations.
- At iteration $j + 1$: As all residual capacities in G_f are integers, then bottleneck $(P, f) \in \mathbb{Z}$, for the augmenting path found in iteration $j + 1$.
- Thus the augmented flow values are integers. □

Max Flow and
Min Cut

Properties of
flows and cuts

Residual
graph

Augmenting
path

MaxFlow
MinCut Thm

Ford
Fulkerson alg

Networks with integer capacities

Max Flow and
Min Cut

Properties of
flows and cuts

Residual
graph

Augmenting
path

MaxFlow
MinCut Thm

Ford
Fulkerson alg

Theorem (**Integrality theorem**)

Let $\mathcal{N} = (V, E, c, s, t)$ where $c : E \rightarrow \mathbb{Z}^+$. There exists a max-flow f^ such that $f^*(e)$ is an integer, for any $e \in E$.*

Proof:

Since the algorithm terminates, the theorem follows from the integrality invariant lemma. □

Networks with integer capacities: FF running time

Max Flow and
Min Cut

Properties of
flows and cuts

Residual
graph

Augmenting
path

MaxFlow
MinCut Thm

Ford
Fulkerson alg

Lemma

*Let C be the min cut capacity (=max. flow value),
Ford-Fulkerson terminates after finding at most C augmenting
paths.*

Proof: The value of the flow increases by ≥ 1 after each
augmentation. □

Networks with integer capacities: FF running time

Max Flow and
Min Cut

Properties of
flows and cuts

Residual
graph

Augmenting
path


MaxFlow
MinCut Thm

Ford
Fulkerson alg

- The number of iterations is $\leq C$. At each iteration:
- Constructing G_f , with $E(G_f) \leq 2m$, takes $O(m)$ time.
- $O(n + m)$ time to find an augmenting path, or deciding that it does not exist.
- Total running time is $O(C(n + m)) = O(Cm)$
- Is that **polynomic**? No, only **pseudopolynomic**

Networks with integer capacities: FF running time


The number of iterations of Ford-Fulkerson could be $\Theta(C)$



Ford-Fulkerson can alternate between the two long paths, and require $2M$ iterations. Taking $M = 10^{10}$, FF on a graph with 4 vertices can take time $2 \cdot 10^{10}$.

Networks with integer capacities: FF running time

The number of iterations of Ford-Fulkerson could be $\Theta(C)$



Ford-Fulkerson can alternate between the two long paths, and require $2M$ iterations. Taking $M = 10^{10}$, FF on a graph with 4 vertices can take time $2 \cdot 10^{10}$.