
Resolution in Propositional Logic

Albert Oliveras and Enric Rodríguez-Carbonell

Logic and Algebra in Computer Science

Session 3

Fall 2009, Barcelona



Overview of the session

- Inference rules
- Resolution
- Ordered resolution
- Practical remarks



Inference rules


- They allow one to deduce new formulas from given ones
- Given an inference rule R and a set of formulas S , we define:
 - The closure of S under R , denoted $R(S)$ is the set of all formulas that can be obtained in zero or more deduction steps from S using R .
 - More formally, for $i \geq 0$

$$\begin{aligned} S_0 &= S \\ S_{i+1} &= S_i \cup R_1(S_i) \end{aligned} \quad \text{and} \quad R(S) = \bigcup_{i=0}^{\infty} S_i$$

where $R_1(S_i)$ is the set of all formula obtained from S_i in exactly one application of R .



Inference rules - Closure



Inference rules - properties

- R is **correct** iff $F \in R(S)$ implies $S \models F$
 - That is, the closure only contains logical consequences (but maybe not all of them)
- R is **complete** iff $S \models F$ implies $F \in R(S)$
 - That is, the closure contains all logical consequences (but maybe something more)
- Ideally, we want **correct** and **complete** inference rules
- A weaker notion of completeness is **refutational completeness**:
$$S \text{ unsatisfiable} \implies \square \in R(S)$$
- If R is correct and refutationally complete, then
$$S \text{ unsatisfiable} \iff \square \in R(S)$$

EXERCISE: prove the last property



Overview of the session

- Inference rules
- **Resolution**
- Ordered resolution
- Practical remarks



Resolution

- The resolution inference rule is the following:

$$\frac{p \vee C \quad \neg p \vee D}{C \vee D}$$

- We will see that:

- Resolution is

- Correct

- Not complete

- Refutationally complete

- If S is a finite set of clauses, then $Res(S)$ is also finite

- Hence, given a set of clauses S , its satisfiability is checked by:

1. Computing $Res(S)$

2. **If** $\square \in Res(S)$ **Then** UNSAT ; **Else** SAT



Resolution - Properties

EXERCISE: prove that

- If S is finite, $Res(S)$ is also finite
- Resolution is not complete
- Resolution is correct
- Resolution is refutationally complete



Overview of the session

- Inference rules
- Resolution
- **Ordered resolution**
- Practical remarks



Ordered resolution

- The proof of refutational completeness introduces **ordered resolution**
- Given clauses S and a total ordering on the variables in S :

$$p_1 < p_2 < p_3 < \dots$$

we can define **ordered resolution**:

$$\frac{p \vee C \quad \neg p \vee D}{C \vee D} \text{ if } p > q \text{ for all var. } q \in C \vee D$$

- It is easy to see that:
 - If S is finite, $ResOrd(S)$ is also finite
 - It is correct (because resolution is)
 - It is refutationally complete (same proof suffices)
- Hence, it is better from the practical point of view



Overview of the session

- Inference rules
- Resolution
- Ordered resolution
- Practical remarks



Practical Remarks

- In practice, even ordered resolution is not efficient enough
- SAT engines based on resolution not used in practice
- However, resolution plays a crucial role in DPLL

